

PROPAGATION OF REGULARITY IN L^p -SPACES FOR KOLMOGOROV TYPE HYPOELLIPTIC OPERATORS

Zhen-Qing Chen and Xicheng Zhang

ABSTRACT. Consider the following Kolmogorov type hypoelliptic operator

$$\mathcal{L} := \sum_{j=2}^n x_j \cdot \nabla_{x_{j-1}} + \Delta_{x_n},$$

where $n \geq 2$ and $x = (x_1, \dots, x_n) \in (\mathbb{R}^d)^n = \mathbb{R}^{nd}$. Let $\{\mathcal{T}_t; t \geq 0\}$ be the semi-group associated with \mathcal{L} . For any $p \in (1, \infty)$, we show that there is a constant $C = C(p, n, d) > 0$ such that for any $f(t, x) \in L^p(\mathbb{R} \times \mathbb{R}^{nd}) = L^p(\mathbb{R}^{1+nd})$,

$$\left\| \Delta_{x_j}^{\frac{1}{1+2(n-j)}} \int_0^\infty \mathcal{T}_t f(t+s, x) dt \right\|_p \leq C \|f\|_p, \quad j = 1, \dots, n,$$

where $\|\cdot\|_p$ is the usual L^p -norm in $L^p(\mathbb{R}^{1+nd}; ds \times dx)$. To show this type of estimates, we first study the propagation of regularity in L^2 -space from variable x_n to x_1 for the solution of the transport equation $\partial_t u + \sum_{j=2}^n x_j \cdot \nabla_{x_{j-1}} u = f$.

1. INTRODUCTION

Let $n \geq 2$ and $d \in \mathbb{N}$. In this paper we consider the following Kolmogorov type hypoelliptic operator:

$$\mathcal{L}_t := \sum_{i,j=1}^d a_t^{ij} \partial_{x_{ni}} \partial_{x_{nj}} + \sum_{j=2}^n x_j \cdot \nabla_{x_{j-1}}, \quad (1.1)$$

where $x = (x_1, x_2, \dots, x_n) \in \mathbb{R}^{nd}$ with $x_j = (x_{j1}, \dots, x_{jd}) \in \mathbb{R}^d$ for each $j = 1, \dots, n$, $\nabla_{x_j} = (\partial_{x_{j1}}, \dots, \partial_{x_{jd}})$, and $a_t = (a_t^{ij}) : \mathbb{R} \rightarrow \mathbb{M}_{sym}^d$ is a measurable map. Here \mathbb{M}_{sym}^d stands for the set of all symmetric $d \times d$ -matrices. Suppose that

$$\kappa^{-1} \mathbb{I}_{d \times d} \leq a_t \leq \kappa \mathbb{I}_{d \times d} \quad (1.2)$$

for some $\kappa \geq 1$. Let $\nabla := (\nabla_{x_1}, \dots, \nabla_{x_n})$, $\nabla_{x_n}^2 := (\partial_{x_{ni}} \partial_{x_{nj}})_{i,j=1, \dots, d}$ and

$$A = A_n = \begin{pmatrix} 0_{d \times d} & \mathbb{I}_{d \times d} & 0_{d \times d} & \cdots & \cdots \\ 0_{d \times d} & 0_{d \times d} & \mathbb{I}_{d \times d} & 0_{d \times d} & \cdots \\ \vdots & \ddots & \ddots & \ddots & \vdots \\ \vdots & \cdots & \ddots & 0_{d \times d} & \mathbb{I}_{d \times d} \\ 0_{d \times d} & \cdots & \cdots & 0_{d \times d} & 0_{d \times d} \end{pmatrix}_{nd \times nd}. \quad (1.3)$$

We can rewrite \mathcal{L}_t as the following compact form:

$$\mathcal{L}_t = \text{tr}(a_t \cdot \nabla_{x_n}^2) + Ax \cdot \nabla,$$

where ‘‘tr’’ denotes the trace of matrix.

Consider the following linear stochastic differential equations (SDEs):

$$dX_{s,t} = AX_{s,t}dt + \sigma_t^a dW_t, \quad X_{s,s} = x, \quad t > s, \quad (1.4)$$

where $(W_t)_{t \in \mathbb{R}}$ is a standard nd -dimensional Brownian motion and

$$\sigma_t^a := \begin{pmatrix} 0_{(n-1)d \times (n-1)d}, & 0_{(n-1)d \times d} \\ 0_{d \times (n-1)d}, & (\sqrt{2a_t})_{d \times d} \end{pmatrix}_{nd \times nd}. \quad (1.5)$$

It is easy to see that the solution $X_{s,t}$ is explicitly given by

$$X_{s,t} := X_{s,t}(x) := e^{(t-s)A}x + \int_s^t e^{(t-r)A} \sigma_r^a dW_r,$$

where e^{tA} is the exponential matrix with the expression

$$e^{tA} = \begin{pmatrix} \mathbb{I}_{d \times d} & t\mathbb{I}_{d \times d} & \frac{t^2 \mathbb{I}_{d \times d}}{2} & \cdots & \frac{t^{n-1} \mathbb{I}_{d \times d}}{(n-1)!} \\ 0_{d \times d} & \mathbb{I}_{d \times d} & t\mathbb{I}_{d \times d} & \cdots & \frac{t^{n-2} \mathbb{I}_{d \times d}}{(n-2)!} \\ \vdots & \ddots & \ddots & \ddots & \vdots \\ \vdots & \cdots & 0_{d \times d} & \mathbb{I}_{d \times d} & t\mathbb{I}_{d \times d} \\ 0_{d \times d} & \cdots & \cdots & 0_{d \times d} & \mathbb{I}_{d \times d} \end{pmatrix}_{nd \times nd}. \quad (1.6)$$

Notice that if $a_t = a$ does not depend on t (i.e., time homogeneous), then

$$X_{s,t}(x) \stackrel{(d)}{=} Z_{t-s}(x) \text{ with } Z_t(x) := e^{tA}x + \int_0^t e^{rA} \sigma^a dW_r.$$

In this case, $Z_t(x)$ is an (nd) -dimensional Gaussian random variable with density

$$p_t(x, y) = \frac{e^{-(\Theta_{t-1/2}(y - e^{tA}x))^* \Sigma^{-1} \Theta_{t-1/2}(y - e^{tA}x)}}{((2\pi)^{nd} t^{n^2d} \det(\Sigma))^{1/2}},$$

where $\Theta_r : \mathbb{R}^{nd} \rightarrow \mathbb{R}^{nd}$ is the dilation operator defined by

$$\Theta_r(x) = (r^{2n-1}x_1, r^{2n-3}x_2, \dots, rx_n), \quad (1.7)$$

and $\Sigma := \int_0^1 e^{rA} \sigma^a (\sigma^a)^* e^{rA*} dr$ is the covariance matrix (see [7] or (2.2) below). For $f \in C_b^2(\mathbb{R}^{nd})$, define

$$\mathcal{T}_{s,t}f(x) := \mathbb{E}f(X_{s,t}(x)). \quad (1.8)$$

It is well-known that \mathcal{L}_t is the infinitesimal generator of $\mathcal{T}_{s,t}$, i.e.,

$$\partial_s \mathcal{T}_{s,t}f + \mathcal{L}_s \mathcal{T}_{s,t}f = 0. \quad (1.9)$$

The goal of this paper is to show the following L^p -maximal regularity estimate.

Theorem 1.1. *Let $p \in (1, \infty)$ and $j = 1, \dots, n$. Under (1.2), there is a constant $C = C(n, \kappa, p, d) > 0$ such that for any $f \in L^p(\mathbb{R}^{1+nd})$ and $\lambda \geq 0$,*

$$\left\| \Delta_{x_j}^{\frac{1}{1+2(n-j)}} \int_0^\infty e^{-\lambda t} \mathcal{T}_{s,t+s}f(t+s, x) dt \right\|_p \leq C \|f\|_p, \quad (1.10)$$

where $\Delta_{x_j}^{\frac{1}{1+2(n-j)}} := -(-\Delta_{x_j})^{\frac{1}{1+2(n-j)}}$ is the fractional Laplacian acting on the variable x_j .

One of the motivation of studying the estimate (1.10) comes from the study of the following $n + 1$ -order stochastic differential equation:

$$dX_t^{(n)} = b_t(X_t, X_t^{(1)}, \dots, X_t^{(n)})dt + \sigma_t(X_t, X_t^{(1)}, \dots, X_t^{(n)})dW_t, \quad (1.11)$$

where $X_t^{(n)}$ denotes the n -order derivative of X_t with respect to the time variable and $b : \mathbb{R}_+ \times \mathbb{R}^{(n+1)d} \rightarrow \mathbb{R}^d$ and $\sigma : \mathbb{R}_+ \times \mathbb{R}^{(n+1)d} \rightarrow \mathbb{R}^d \otimes \mathbb{R}^d$ are two measurable functions. Notice that if we let

$$\mathbf{X}_t := (X_t, X_t^{(1)}, \dots, X_t^{(n)}),$$

then \mathbf{X}_t solves the following one order stochastic differential equation

$$d\mathbf{X}_t = (X_t^{(1)}, \dots, X_t^{(n)}, b_t(\mathbf{X}_t))dt + (0, \dots, 0, \sigma_t(\mathbf{X}_t)dW_t), \quad \mathbf{X}_0 = \mathbf{x},$$

where $\mathbf{x} = (\mathbf{x}_i)_{i=0, \dots, n} = ((x_{ij})_{j=1, \dots, d})_{i=0, \dots, n}$. In particular, the infinitesimal generator of Markov process $\mathbf{X}_t(\mathbf{x})$ is given by

$$\mathcal{L}_t f(\mathbf{x}) = \sum_{i,j,k=1}^d (\sigma_t^{ik} \sigma_t^{jk})(\mathbf{x}) \partial_{x_{ni}} \partial_{x_{nj}} f(\mathbf{x}) + \sum_{j=1}^n \mathbf{x}_j \cdot \nabla_{\mathbf{x}_{j-1}} f(\mathbf{x}) + b_t(\mathbf{x}) \cdot \nabla_{\mathbf{x}_n} f(\mathbf{x}),$$

which takes the same form as (1.1). Thus, the estimate (1.10) could be used to study the well-posedness of SDE (1.11) with rough coefficients b and σ . Indeed, when $n = 1$ and σ is bounded and uniformly nondegenerate, the second named author [10] studied the strong well-posedness of SDE (1.11) with $(\mathbb{I} - \Delta_{\mathbf{x}_1})^{1/3} b, \nabla \sigma \in L_{loc}^p(\mathbb{R}_+ \times \mathbb{R}^{2d})$ for some $p > 4d + 2$. See also [5] for similar results when $\sigma = \mathbb{I}$.

Let us now recall some related results in literature about the estimate (1.10). In [3], the authors adopted Coifman-Weiss' theorem to show the estimate (1.10) for $j = n$. When $n = 2$, in [4], we used Fefferman-Stein's theorem to show the estimate (1.10) for $j = 1, 2$ even for nonlocal operators. It should be noticed that the methods used in [3] and [4] are quite different. In [3], the key point is to show some weak 1-1 type estimate. While in [4], the main point is to show that the operator in (1.10) is bounded from L^∞ to some BMO spaces. In particular, to show the propagation of the regularity from the nondegenerate component to the degenerate component, in [4], we have used Bouchet's result [2]. More precisely, Bouchet studied the following transport equation:

$$\partial_t u + x_2 \cdot \nabla_{x_1} u = f,$$

and showed that for any $\alpha \geq 0$,

$$\|\Delta_{x_1}^{\frac{\alpha}{2(1+\alpha)}} u\|_2 \leq C \|\Delta_{x_2}^{\frac{\alpha}{2}} u\|_2^{\frac{1}{1+\alpha}} \|f\|_2^{\frac{\alpha}{1+\alpha}},$$

where $C = C(\alpha, d) > 0$. A simplified proof of this type estimate was provided in [1]. Thus, the first goal of this paper is to extend the above estimate to the following more general transport equation:

$$\partial_t u + \sum_{j=2}^n x_j \cdot \nabla_{x_{j-1}} u = f.$$

That is, we want to show that for any $j = 1, \dots, n-1$ and $\alpha \geq 0$, there is a constant $C = C(\alpha, d, j, n) > 0$ such that

$$\|\Delta_{x_j}^{\frac{\alpha}{2(1+\alpha)}} u\|_2 \leq C \|\Delta_{x_{j+1}}^{\frac{\alpha}{2}} u\|_2^{\frac{1}{1+\alpha}} \|f\|_2^{\frac{\alpha}{1+\alpha}}.$$

It seems that this type estimate is not trivial compared with [1], see Section 3.

Although the above result is proven for Laplacian operator, we can extend it to more general nonlocal operator as in [4] without any difficulty. Indeed, let us consider the following nonlocal operator:

$$\widetilde{\mathcal{L}}_\sigma^\nu f(x) := \int_{\mathbb{R}^d} [f(x + \sigma y) + f(x - y) - 2f(x)] \nu(dy),$$

where $\sigma \in \mathbb{M}^d$ is a $d \times d$ matrix and ν is a symmetric Lévy measure. Let $n \geq 2$ and

$$\mathcal{L}_t f(x) := \widetilde{\mathcal{L}}_{\sigma_t, x_n}^{\nu_t} f(x) + \sum_{j=2}^n x_j \cdot \nabla_{x_{j-1}} f(x),$$

where $\widetilde{\mathcal{L}}_{\sigma_t, x_n}^{\nu_t}$ means that the operator acts on the variable x_n . Suppose that

$$\|\sigma\|_\infty + \|\sigma^{-1}\|_\infty < \infty$$

and for some $\alpha \in (0, 2)$,

$$\nu_1^{(\alpha)} \leq \nu_s \leq \nu_2^{(\alpha)},$$

where $\nu_1^{(\alpha)}$ and $\nu_2^{(\alpha)}$ are two symmetric and nondegenerate α -stable Lévy measures (see [4]). Under the above assumptions, as in [4], we can show that for any $j = 1, \dots, n$,

$$\left\| \Delta_{x_j}^{\frac{\alpha}{2(1+\alpha(n-j))}} \int_0^\infty e^{-\lambda t} \mathcal{T}_{s,t+s}^{\nu, \sigma} f(t+s, x) dt \right\|_p \leq C \|f\|_p, \quad (1.12)$$

where $\mathcal{T}_{s,t}^{\nu, \sigma}$ is defined as in (1.8) by using the non-stationary Lévy process L_t with Lévy measure ν_t to replace the Brownian motion. We note that at the almost same time, Huang, Menozzi and Priola [6] have obtained (1.12) for time-independent σ and ν by using Coifman-Weiss' theorem. As mentioned above, our proof is based on Fefferman-Stein's theorem.

This paper is organized as follows: In Section 2, we prepare some estimates about the distribution density of $X_{s,t}(x)$, and give the necessary Fefferman-Stein's theorem. In Section 3, we show the propagation of the regularity for transport equation. In Section 4, we prove our main result.

Throughout this paper, we use the following convention: The letters C and c with or without subscripts will denote a positive constant, whose value may change in different places. Moreover, we use $A \lesssim B$ to denote $A \leq CB$ for some constant $C > 0$.

2. PRELIMINARIES

2.1. Estimate of density of $X_{s,t}(x)$. In this subsection we show some necessary estimate about the density of $X_{s,t}(x)$.

Lemma 2.1. Under (1.2), $X_{s,t}(0)$ admits a smooth density $p_{s,t}^{(0)}(y)$ so that for each $\beta = (\beta_1, \dots, \beta_n) \in \mathbb{N}_0^n$, where $\mathbb{N}_0 = \{0\} \cup \mathbb{N}$, there are constants $C, c > 0$ only depending on n, β, d and κ such that for all $s < t$ and $x, y \in \mathbb{R}^{nd}$,

$$|\nabla_{y_1}^{\beta_1} \dots \nabla_{y_n}^{\beta_n} p_{s,t}^{(0)}(y)| \leq C(t-s)^{-(n^2d + \sum_{i=1}^n (2(n-i)+1)\beta_i)/2} e^{-c|\Theta_{(t-s)^{-1/2}y}|^2}, \quad (2.1)$$

where Θ_r is the dilation operator defined by (1.7).

Proof. Since $(cW_{\cdot/c^2}) \stackrel{(d)}{=} W$ for $c \neq 0$, by the change of variables, we have

$$X_{s,t}(0) = \int_s^t e^{(t-r)A} \sigma_r^a dW_r \stackrel{(d)}{=} (t-s)^{1/2} \int_0^1 e^{(t-s)(1-r)A} \sigma_{s+(t-s)r}^a dW_r.$$

Hence, by definitions (1.5), (1.6) and (1.7),

$$\Theta_{(t-s)^{-1/2}} X_{s,t}(0) \stackrel{(d)}{=} \int_0^1 e^{(1-r)A} \sigma_{s+(t-s)r}^a dW_r \stackrel{(d)}{=} \int_0^1 e^{rA} \tilde{\sigma}_r^a dW_r =: Z,$$

where $\tilde{\sigma}_r^a := \sigma_{s+(t-s)(1-r)}^a$. Since Z is a nd -dimensional Gaussian random variable with mean value zero and covariance matrix

$$\Sigma = \int_0^1 e^{rA} \tilde{\sigma}_r^a (\tilde{\sigma}_r^a)^* e^{rA^*} dr,$$

we have

$$p_{s,t}^{(0)}(y) = \frac{e^{-(\Theta_{(t-s)^{-1/2}y})^* \Sigma^{-1} (\Theta_{(t-s)^{-1/2}y})}}{((2\pi)^{nd} (t-s)^{n^2d} \det(\Sigma))^{1/2}}. \quad (2.2)$$

On the other hand, by (1.5), (1.6) and (1.3), we have for all $y \in \mathbb{R}^{nd}$,

$$\begin{aligned} y^* \Sigma y &= \int_0^1 |y^* e^{rA} \tilde{\sigma}_r^a|^2 dr \geq 2\kappa^{-1} \int_0^1 \left| \sum_{j=1}^n \frac{r^{n-j}}{(n-j)!} y_j \right|^2 dr \\ &\geq 2\kappa^{-1} |y|^2 \inf_{|\omega|=1} \int_0^1 \left| \sum_{j=1}^n \frac{r^{n-j}}{(n-j)!} \omega_j \right|^2 dr. \end{aligned}$$

Since the unit sphere in \mathbb{R}^{nd} is compact, and for each $\mathbb{R}^{nd} \ni \omega \neq 0$,

$$\delta(\omega) := \int_0^1 \left| \sum_{j=1}^n \frac{r^{n-j}}{(n-j)!} \omega_j \right|^2 dr > 0,$$

we obtain that for some $c_0 = c_0(n, \kappa, d) > 0$ and all $y \in \mathbb{R}^{nd}$,

$$y^* \Sigma y \geq 2\kappa^{-1} |y|^2 \inf_{|\omega|=1} \delta(\omega) \Rightarrow y^* \Sigma^{-1} y \geq c_0 |y|^2. \quad (2.3)$$

The desired estimate now follows by the chain rule, (2.2) and (2.3). \square

For $\alpha \in (0, 2]$, the fractional Laplacian $\Delta^{\alpha/2}$ in \mathbb{R}^d is defined by Fourier's transformation as

$$\widehat{\Delta^{\alpha/2} f}(\xi) = |\xi|^\alpha \hat{f}(\xi), \quad f \in \mathcal{S}(\mathbb{R}^d),$$

where $\mathcal{S}(\mathbb{R}^d)$ is the space of Schwartz rapidly decreasing functions. For $\alpha \in (0, 2)$, up to a multiplying constant, an alternative definition of $\Delta^{\alpha/2}$ is given by the following integral form (cf. [8]):

$$\Delta^{\alpha/2} f(x) = \int_{\mathbb{R}^d} \frac{\delta_z^{(2)} f(x)}{|z|^{d+\alpha}} dz, \quad (2.4)$$

where $\delta_z^{(2)} f(x) := f(x+z) + f(x-z) - 2f(x)$. Observe that for $\alpha \in (0, 1)$,

$$\Delta^{\alpha/2} f(x) = 2 \int_{\mathbb{R}^d} \frac{\delta_z^{(1)} f(x)}{|z|^{d+\alpha}} dz \text{ with } \delta_z^{(1)} f(x) := f(x+z) - f(x). \quad (2.5)$$

Corollary 2.2. *For any $j = 1, \dots, n$, $\alpha \in (0, 2]$ and $\beta = (\beta_1, \dots, \beta_n) \in \mathbb{N}_0^n$, there is a constant $C > 0$ such that for all $f \in C_b^\infty(\mathbb{R}^{nd})$ and $s < t$,*

$$\|\Delta_{x_j}^{\alpha/2} \nabla_{x_1}^{\beta_1} \dots \nabla_{x_n}^{\beta_n} \mathcal{T}_{s,t} f\|_\infty \leq C |t-s|^{-\left(\sum_{i=1}^n (2(n-i)+1)\beta_i + (2(n-j)+1)\alpha\right)/2} \|f\|_\infty, \quad (2.6)$$

$$\|\nabla_{x_1}^{\beta_1} \dots \nabla_{x_n}^{\beta_n} \mathcal{T}_{s,t} \Delta_{x_j}^{\alpha/2} f\|_\infty \leq C |t-s|^{-\left(\sum_{i=1}^n (2(n-i)+1)\beta_i + (2(n-j)+1)\alpha\right)/2} \|f\|_\infty, \quad (2.7)$$

where $\Delta_{x_j}^{\alpha/2}$ means that the fractional Laplacian acts on the variable x_j .

Proof. Below we only show (2.6) and (2.7) for $\alpha \in (0, 2)$. Let $p_{s,t}(x, y)$ be the distribution density of $X_{s,t}(x) = e^{(t-s)A}x + X_{s,t}(0)$. We have

$$p_{s,t}(x, y) = p_{s,t}^{(0)}(y - e^{(t-s)A}x). \quad (2.8)$$

For simplicity of notation, we write

$$h_y(x) := \nabla_{x_1}^{\beta_1} \dots \nabla_{x_n}^{\beta_n} p_{s,t}(x, y), \quad \gamma := \sum_{i=1}^n (2(n-i)+1)\beta_i/2$$

and

$$\delta_{z_j}^{(2)} h_y(x) = h_y(x + \tilde{z}_j) + h_y(x - \tilde{z}_j) - 2h_y(x), \quad \tilde{z}_j = (0, \dots, z_j, \dots, 0).$$

By (2.1), (2.8) and the chain rule, we have

$$\begin{aligned} |\delta_{z_j}^{(2)} h_y(x)| &\leq C(t-s)^{-n^2 d/2 - \gamma} \left(e^{-c|\Theta_{(t-s)^{-1/2}}(y - e^{(t-s)A}(x + \tilde{z}_j))|^2} \right. \\ &\quad \left. + e^{-c|\Theta_{(t-s)^{-1/2}}(y - e^{(t-s)A}(x - \tilde{z}_j))|^2} + e^{-c|\Theta_{(t-s)^{-1/2}}(y - e^{(t-s)A}x)|^2} \right), \end{aligned} \quad (2.9)$$

and also by the mean value formula,

$$|\delta_{z_j}^{(2)} h_y(x)| \leq C(t-s)^{-n^2 d/2 - \gamma - (2(n-j)+1)} e^{-c|\Theta_{(t-s)^{-1/2}}(y - e^{(t-s)A}\tilde{x})|^2} |z_j|^2 \quad (2.10)$$

for some $\tilde{x} \in \mathbb{R}^{nd}$ depending on z_j . By formula (2.4), we have

$$\begin{aligned} \Delta_{x_j}^{\alpha/2} h_y(x) &= \left(\int_{|z_j| > (t-s)^{(2(n-j)+1)/2}} + \int_{|z_j| \leq (t-s)^{(2(n-j)+1)/2}} \right) \frac{\delta_{z_j}^{(2)} h_y(x)}{|z_j|^{d+\alpha}} dz_j \\ &=: I_1(x, y) + I_2(x, y). \end{aligned}$$

For $I_1(x, y)$, by (2.9) we have

$$\int_{\mathbb{R}^{nd}} |I_1(x, y)| dy \lesssim (t-s)^{-\gamma} \int_{|z_j| > (t-s)^{(2(n-j)+1)/2}} \frac{dz_j}{|z_j|^{d+\alpha}} \lesssim (t-s)^{-\gamma - (2(n-j)+1)\alpha/2},$$

where we have used that

$$(t-s)^{-n^2 d/2} \int_{\mathbb{R}^{nd}} e^{-c|\Theta_{(t-s)^{-1/2}(y)}|^2} dy = \int_{\mathbb{R}^{nd}} e^{-c|y|^2} dy.$$

For $I_2(x, y)$, by (2.10) we have

$$\begin{aligned} \int_{\mathbb{R}^{nd}} |I_2(x, y)| dy &\lesssim (t-s)^{-\gamma-(2(n-j)+1)} \int_{|z_j| \leq (t-s)^{(2(n-j)+1)/2}} \frac{|z_j|^2}{|z_j|^{d+\alpha}} dz_j \\ &\lesssim (t-s)^{-\gamma-(2(n-j)+1)\alpha/2}. \end{aligned}$$

Combining the above calculations, we obtain

$$|\Delta_{x_j}^{\alpha/2} \nabla_{x_1}^{\beta_1} \cdots \nabla_{x_n}^{\beta_n} \mathcal{T}_{s,t} f(x)| = \left| \int_{\mathbb{R}^d} \Delta_{x_j}^{\alpha/2} h_y(x) f(y) dy \right| \leq C(t-s)^{-\gamma-(2(n-j)+1)\alpha/2} \|f\|_{\infty}.$$

Thus we proved (2.6). Similarly, we can show (2.6). \square

2.2. Fefferman-Stein's theorem. First of all, we introduce a family of “balls” in \mathbb{R}^{1+nd} . For any $r > 0$ and point $(t_0, x_0) \in \mathbb{R}^{1+nd}$, we define

$$Q_r(t_0, x_0) := \{(t, x) : \ell(t - t_0, x - e^{(t-t_0)A} x_0) \leq r\},$$

where

$$\ell(t, x) := \max\left(|t|^{\frac{1}{2}}, |x_1|^{\frac{1}{2n-1}}, |x_2|^{\frac{1}{2n-3}}, \dots, |x_{n-1}|^{\frac{1}{3}}, |x_n|\right).$$

The set of all such balls is denoted by \mathbb{Q} . We have

Lemma 2.3. (i) $\ell(r^2 t, \Theta_r x) = r \ell(t, x)$ for any $r > 0$, where Θ_r is the dilation operator defined by (1.7).

(ii) $|Q_r(t_0, x_0)| = \omega_d^n r^{n^2 d + 2}$, where $|\cdot|$ denotes the Euclidean volume and ω_d is the volume of the unit ball in \mathbb{R}^d .

(iii) For all $(t, x), (s, y), (r, z) \in \mathbb{R}^{1+nd}$, we have

$$\begin{aligned} \ell(s-t, y - e^{(s-t)A} x) &\leq 3\ell(t-s, x - e^{(t-s)A} y) \leq \\ &\leq 12\left(\ell(t-r, x - e^{(t-r)A} z) + \ell(r-s, z - e^{(r-s)A} y)\right). \end{aligned} \tag{2.11}$$

(iv) Suppose that $Q_r(t_0, x_0) \cap Q_r(t'_0, x'_0) \neq \emptyset$, then

$$Q_r(t_0, x_0) \subset Q_{20r}(t'_0, x'_0). \tag{2.12}$$

(v) For $(t, x), (s, y) \in \mathbb{R}^{1+nd}$, define

$$\rho((t, x), (s, y)) := \ell(t-s, x - e^{(t-s)A} y) + \ell(s-t, y - e^{(s-t)A} x),$$

and for $(t_0, x_0) \in \mathbb{R}^{1+nd}$ and $r > 0$,

$$\tilde{Q}_r(t_0, x_0) := \{(t, x) : \rho((t, x), (t_0, x_0)) \leq r\}.$$

Then $Q_r(t_0, x_0) \subset \tilde{Q}_r(t_0, x_0) \subset Q_{4r}(t_0, x_0)$.

Proof. (i) and (ii) are direct by definition.

(iii) We only prove the second inequality in (2.11). The first one is similar. Observing that for all $(t, x), (s, y) \in \mathbb{R}^{1+nd}$,

$$\ell(t + s, x + y) \leq \ell(t, x) + \ell(s, y), \quad (2.13)$$

we have

$$\ell(t - s, x - e^{(t-s)A}y) \leq \ell(t - r, x - e^{(t-r)A}z) + \ell(r - s, e^{(t-r)A}z - e^{(t-s)A}y).$$

For simplicity, we write

$$a := \ell(t - r, x - e^{(t-r)A}z), \quad b := \ell(r - s, z - e^{(r-s)A}y).$$

By the definition of ℓ , we have

$$|t - r| \leq a^2, \quad |(z - e^{(r-s)A}y)_j| \leq b^{1+2(n-j)}, \quad j = 1, \dots, n.$$

Hence, for each $i = 1, \dots, n$,

$$\begin{aligned} |(e^{(t-r)A}z - e^{(t-s)A}y)_i| &= \left| \sum_{j=1}^n (e^{(t-r)A})_{ij} (z - e^{(r-s)A}y)_j \right| \\ &\stackrel{(1.6)}{\leq} \sum_{j=i}^n \frac{|t - r|^{j-i}}{(j-i)!} b^{1+2(n-j)} \leq \sum_{j=i}^n \frac{a^{2(j-i)}}{(j-i)!} b^{1+2(n-j)} \\ &\leq (a \vee b)^{1+2(n-i)} \sum_{j=i}^n \frac{1}{(j-i)!} \leq 3(a \vee b)^{1+2(n-i)}, \end{aligned} \quad (2.14)$$

and

$$\ell(t - s, x - e^{(t-s)A}y) \leq a + 3(a \vee b) \leq 4(a + b).$$

(iv) and (v) are easy consequences of (iii). \square

Now, for any $f \in L^1_{loc}(\mathbb{R}^{1+nd})$, we define the Hardy-Littlewood maximal function by

$$\mathcal{M}f(t, x) := \sup_{r>0} \int_{Q_r(t,x)} |f(t', x')| dx' dt',$$

and the sharp function by

$$\mathcal{M}^\sharp f(t, x) := \sup_{r>0} \int_{Q_r(t,x)} |f(t', x') - f_{Q_r(t,x)}| dx' dt',$$

where for a $Q \in \mathbb{Q}$,

$$f_Q := \int_Q f(t', x') dx' dt' = \frac{1}{|Q|} \int_Q f(t', x') dx' dt'.$$

One says that a function $f \in BMO(\mathbb{R}^{1+nd})$ if $\mathcal{M}^\sharp f \in L^\infty(\mathbb{R}^{1+nd})$. Clearly, $f \in BMO(\mathbb{R}^{1+nd})$ if and only if there exists a constant $C > 0$ such that for any $Q \in \mathbb{Q}$, and for some $c_Q \in \mathbb{R}$,

$$\int_Q |f(t', x') - c_Q| dx' dt' \leq C.$$

The following version of Fefferman-Stein's theorem can be proven as in [4, Theorem 2.12] by Lemma 2.3. We omit the details.

Theorem 2.4. *For $q \in (1, \infty)$, let \mathcal{P} be a bounded linear operator from $L^q(\mathbb{R}^{1+nd})$ to $L^q(\mathbb{R}^{1+nd})$ and also from $L^\infty(\mathbb{R}^{1+nd})$ to $BMO(\mathbb{R}^{1+nd})$. Then for any $p \in [q, \infty)$ and $f \in L^p(\mathbb{R}^{1+nd})$,*

$$\|\mathcal{P}f\|_p \leq C\|f\|_p,$$

where the constant C depends only on p, q and the norms of $\|\mathcal{P}\|_{L^q \rightarrow L^q}$ and $\|\mathcal{P}\|_{L^\infty \rightarrow BMO}$.

3. PROPAGATION OF REGULARITY IN L^2 -SPACE FOR TRANSPORT EQUATIONS

Fix $n \geq 2$ and $\lambda \geq 0$. Let A be as in (1.3). In this section we consider the following linear transport equation in \mathbb{R}^{nd} :

$$\partial_s u + Ax \cdot \nabla u - \lambda u + f = 0. \quad (3.1)$$

Taking Fourier's transformation in the spatial variable x , we obtain

$$\partial_s \hat{u} - A^* \xi \cdot \nabla \hat{u} - \lambda \hat{u} + \hat{f} = 0, \quad (3.2)$$

where A^* is the transpose of A , and ξ is the dual variable. Multiplying both sides by the complex conjugate of \hat{u} , we get

$$\partial_s |\hat{u}|^2 - A^* \xi \cdot \nabla |\hat{u}|^2 - \lambda |\hat{u}|^2 + 2\text{Re}(\hat{f}, \bar{\hat{u}}) = 0.$$

Let $\phi(\xi)$ be a smooth function and define

$$g_\phi := 2\phi \text{Re}(\hat{f}, \bar{\hat{u}}) + A^* \xi \cdot \nabla \phi |\hat{u}|^2. \quad (3.3)$$

We have

$$\partial_s (|\hat{u}|^2 \phi) - A^* \xi \cdot \nabla (|\hat{u}|^2 \phi) - \lambda (|\hat{u}|^2 \phi) + g_\phi = 0, \quad (3.4)$$

and if \hat{u} has compact support, then

$$\begin{aligned} (|\hat{u}|^2 \phi)(s, \xi) &= - \int_0^\infty \partial_t (e^{-\lambda t} (|\hat{u}|^2 \phi)(t + s, e^{-tA^*} \xi)) dt \\ &= \int_0^\infty e^{-\lambda t} g_\phi(t + s, e^{-tA^*} \xi) dt, \end{aligned} \quad (3.5)$$

where $e^{-tA^*} = (e^{-tA})^*$ is the transpose of exponential matrix in (1.6).

The following propagation of regularity in L^2 -space is the key step in the proof of Theorem 1.1.

Theorem 3.1. *Let $f, u \in L^2(\mathbb{R}^{1+nd})$ so that (3.1) holds in the weak sense. For any $\alpha \geq 0$ and $j = 1, 2, \dots, n-1$, there is a constant $C = C(\alpha, j, d) > 0$ such that*

$$\|\Delta_{x_j}^{\frac{\alpha}{2(1+\alpha)}} u\|_2 \leq C \|\Delta_{x_{j+1}}^{\frac{\alpha}{2}} u\|_2^{\frac{1}{1+\alpha}} \|f\|_2^{\frac{\alpha}{1+\alpha}}. \quad (3.6)$$

In particular,

$$\|\Delta_{x_j}^{\frac{\alpha}{2(1+(n-j)\alpha)}} u\|_2 \leq C \|\Delta_{x_n}^{\frac{\alpha}{2}} u\|_2^{\frac{1+(n-j-1)\alpha}{1+(n-j)\alpha}} \|f\|_2^{\frac{\alpha}{1+(n-j)\alpha}}. \quad (3.7)$$

Proof. Let $\rho : [0, \infty) \rightarrow [0, 1]$ be a smooth function with $\rho(s) = 1$ for $s < 1$ and $\rho(s) = 0$ for $s > 2$. For $R > 0$, let $\chi_R(s, \xi) := \rho(s/R)\rho(|\xi|/R)$ and

$$\hat{u}_R := \chi_R \hat{u}, \quad \hat{f}_R := (A^* \xi \cdot \nabla \chi_R - \partial_s \chi_R) \hat{u} + \chi_R \hat{f}.$$

Since \hat{u}_R satisfies

$$\partial_s \hat{u}_R - A^* \xi \cdot \nabla \hat{u}_R - \lambda \hat{u}_R + \hat{f}_R = 0,$$

if we can show that for some $C = C(\alpha, j, d) > 0$,

$$\|\Delta_{x_j}^{\frac{\alpha}{2(1+\alpha)}} u_R\|_2 \leq C \|\Delta_{x_{j+1}}^{\frac{\alpha}{2}} u_R\|_2^{\frac{1}{1+\alpha}} \|f_R\|_2^{\frac{\alpha}{1+\alpha}},$$

then letting $R \rightarrow \infty$, we get (3.6). Hence, without loss of generality, in the following, we may and shall assume that \hat{u} has compact support. We use the induction method.

Let us first look at the case of $j = 1$. We follow the simple argument of Alexander [1]. For any $\varepsilon > 0$, by Planchel's identity, we have

$$\begin{aligned} \|\Delta_{x_1}^{\frac{\alpha}{2(1+\alpha)}} u\|_2^2 &= \int_{-\infty}^{\infty} \int_{\mathbb{R}^{nd}} |\xi_1|^{\frac{2\alpha}{1+\alpha}} |\hat{u}(s, \xi)|^2 d\xi ds \\ &= \int_{-\infty}^{\infty} \int_{\mathbb{R}^{nd}} |\xi_1|^{\frac{2\alpha}{1+\alpha}} \mathbf{1}_{\{\varepsilon|\xi_2| > |\xi_1|^{1/(1+\alpha)}\}} |\hat{u}(s, \xi)|^2 d\xi ds \\ &\quad + \int_{-\infty}^{\infty} \int_{\mathbb{R}^{nd}} |\xi_1|^{\frac{2\alpha}{1+\alpha}} \mathbf{1}_{\{\varepsilon|\xi_2| \leq |\xi_1|^{1/(1+\alpha)}\}} |\hat{u}(s, \xi)|^2 d\xi ds \\ &=: I_1(\varepsilon) + I_2(\varepsilon). \end{aligned}$$

For $I_1(\varepsilon)$, we have

$$I_1(\varepsilon) \leq \varepsilon^{2\alpha} \int_{-\infty}^{\infty} \int_{\mathbb{R}^{nd}} |\xi_2|^{2\alpha} |\hat{u}(s, \xi)|^2 d\xi ds = \varepsilon^{2\alpha} \|\Delta_{x_2}^{\frac{\alpha}{2}} u\|_2^2.$$

For $I_2(\varepsilon)$, letting $\Omega := \{\varepsilon|\xi_2| \leq |\xi_1|^{1/(1+\alpha)}\}$ and by (3.5) with $\phi = 1$, we have

$$\begin{aligned} I_2(\varepsilon) &= 2 \int_{-\infty}^{\infty} \int_{\mathbb{R}^{nd}} |\xi_1|^{\frac{2\alpha}{1+\alpha}} \mathbf{1}_{\Omega}(\xi) \int_0^{\infty} e^{-\lambda t} \operatorname{Re}(\hat{f}, \bar{\hat{u}})(t + s, e^{-tA^*} \xi) dt d\xi ds \\ &= 2 \int_{-\infty}^{\infty} \int_{\mathbb{R}^{nd}} \int_0^{\infty} |(e^{tA^*} \xi)_1|^{\frac{2\alpha}{1+\alpha}} \mathbf{1}_{\Omega}(e^{tA^*} \xi) e^{-\lambda t} \operatorname{Re}(\hat{f}, \bar{\hat{u}})(t + s, \xi) dt d\xi ds \\ &= 2 \int_{-\infty}^{\infty} \int_{\mathbb{R}^{nd}} \left(\int_0^{\infty} |(e^{tA^*} \xi)_1|^{\frac{2\alpha}{1+\alpha}} \mathbf{1}_{\Omega}(e^{tA^*} \xi) e^{-\lambda t} dt \right) \operatorname{Re}(\hat{f}, \bar{\hat{u}})(s, \xi) ds d\xi. \end{aligned}$$

Observing that

$$e^{tA^*} \xi = \left(\xi_1, t\xi_1 + \xi_2, \frac{t^2}{2}\xi_1 + t\xi_2 + \xi_3, \dots, \frac{t^{n-1}}{(n-1)!}\xi_1 + \dots + t\xi_{n-1} + \xi_n \right),$$

since $\lambda \geq 0$, we have

$$\begin{aligned} \int_0^{\infty} |(e^{tA^*} \xi)_1|^{\frac{2\alpha}{1+\alpha}} \mathbf{1}_{\Omega}(e^{tA^*} \xi) e^{-\lambda t} dt &= |\xi_1|^{\frac{2\alpha}{1+\alpha}} \int_0^{\infty} \mathbf{1}_{\{\varepsilon|\xi_2 + t\xi_1| \leq |\xi_1|^{1/(1+\alpha)}\}} e^{-\lambda t} dt \\ &\leq |\xi_1|^{\frac{2\alpha}{1+\alpha}} \int_{-\infty}^{\infty} \mathbf{1}_{\{|\xi_2|/|\xi_1| - |\xi_1|^{-\alpha/(1+\alpha)}/\varepsilon \leq t \leq |\xi_2|/|\xi_1| + |\xi_1|^{-\alpha/(1+\alpha)}/\varepsilon\}} dt = 2|\xi_1|^{\frac{\alpha}{1+\alpha}}/\varepsilon. \end{aligned} \tag{3.8}$$

Hence, by Young's inequality we have

$$\begin{aligned}
I_2(\varepsilon) &\leq \frac{4}{\varepsilon} \int_{-\infty}^{\infty} \int_{\mathbb{R}^{nd}} |\xi_1|^{\frac{\alpha}{1+\alpha}} (|\hat{f}| |\hat{u}|)(s, \xi) \, ds \, d\xi \\
&\leq \frac{1}{2} \int_{-\infty}^{\infty} \int_{\mathbb{R}^{nd}} |\xi_1|^{\frac{2\alpha}{1+\alpha}} |\hat{u}|^2(s, \xi) \, ds \, d\xi + \frac{8}{\varepsilon^2} \int_{-\infty}^{\infty} \int_{\mathbb{R}^{nd}} |\hat{f}|^2(s, \xi) \, ds \, d\xi \\
&= \frac{1}{2} \|\Delta_{x_1}^{\frac{\alpha}{2(1+\alpha)}} u\|_2^2 + \frac{8}{\varepsilon^2} \|f\|_2^2.
\end{aligned}$$

Combining the above calculations, we obtain

$$\|\Delta_{x_1}^{\frac{\alpha}{2(1+\alpha)}} u\|_2^2 \leq \varepsilon^{2\alpha} \|\Delta_{x_2}^{\frac{\alpha}{2}} u\|_2^2 + \frac{1}{2} \|\Delta_{x_1}^{\frac{\alpha}{2(1+\alpha)}} u\|_2^2 + \frac{8}{\varepsilon^2} \|f\|_2^2,$$

which in turn gives (3.6) for $j = 1$ by letting $\varepsilon = (\|f\|_2 / \|\Delta_{x_2}^{\frac{\alpha}{2}} u\|_2)^{\frac{1}{1+\alpha}}$.

Suppose now that (3.6) has been proven for $j = 1, \dots, k$ with $k \leq n - 2$. We want to show that (3.6) holds for $j = k + 1$. For $\delta > 0$, we define

$$\phi_\delta(\xi) := \prod_{j=1}^k \chi\left(|\xi_j| / (\delta^{k-j+1} |\xi_{k+1}|^{(1+(k-j+2)\alpha)/(1+\alpha)})\right),$$

and write

$$\begin{aligned}
\|\Delta_{x_{k+1}}^{\frac{\alpha}{2(1+\alpha)}} u\|_2 &= \int_{-\infty}^{\infty} \int_{\mathbb{R}^{nd}} |\xi_{k+1}|^{\frac{2\alpha}{1+\alpha}} |\hat{u}(s, \xi)|^2 \, d\xi \, ds \\
&= \int_{-\infty}^{\infty} \int_{\mathbb{R}^{nd}} |\xi_{k+1}|^{\frac{2\alpha}{1+\alpha}} (1 - \phi_\delta(\xi)) |\hat{u}(s, \xi)|^2 \, d\xi \, ds \\
&\quad + \int_{-\infty}^{\infty} \int_{\mathbb{R}^{nd}} |\xi_{k+1}|^{\frac{2\alpha}{1+\alpha}} \phi_\delta(\xi) |\hat{u}(s, \xi)|^2 \, d\xi \, ds \\
&=: K_1(\delta) + K_2(\delta).
\end{aligned}$$

We first treat $K_1(\delta)$. By the induction hypothesis, one sees that for $j = 1, 2, \dots, k$,

$$\|\Delta_{x_j}^{\frac{\alpha}{2(1+(k-j+2)\alpha)}} u\|_2 \leq C \|\Delta_{x_{k+1}}^{\frac{\alpha}{2(1+\alpha)}} u\|_2^{\frac{1+\alpha}{1+(k-j+2)\alpha}} \|f\|_2^{\frac{(k-j+1)\alpha}{1+(k-j+2)\alpha}}. \quad (3.9)$$

Observing that

$$1 - \phi_\delta(\xi) \leq \sum_{j=1}^k \mathbf{1}_{\{|\xi_j| \geq \delta^{k-j+1} |\xi_{k+1}|^{(1+(k-j+2)\alpha)/(1+\alpha)}\}},$$

by (3.9) and Young's inequality, we have

$$\begin{aligned}
K_1(\delta) &\leq \sum_{j=1}^k \int_{-\infty}^{\infty} \int_{\mathbb{R}^{nd}} |\xi_{k+1}|^{\frac{2\alpha}{1+\alpha}} \mathbf{1}_{\{|\xi_j| \geq \delta^{k-j+1} |\xi_{k+1}|^{(1+(k-j+2)\alpha)/(1+\alpha)}\}} |\hat{u}(s, \xi)|^2 \, d\xi \, ds \\
&\leq \sum_{j=1}^k \delta^{-\frac{2(k-j+1)\alpha}{1+(k-j+2)\alpha}} \int_{-\infty}^{\infty} \int_{\mathbb{R}^{nd}} |\xi_j|^{\frac{2\alpha}{1+(k-j+2)\alpha}} |\hat{u}(s, \xi)|^2 \, d\xi \, ds \\
&= \sum_{j=1}^k \delta^{-\frac{2(k-j+1)\alpha}{1+(k-j+2)\alpha}} \|\Delta_{x_j}^{\frac{\alpha}{2(1+(k-j+2)\alpha)}} u\|_2^2
\end{aligned}$$

$$\begin{aligned}
&\leq C \sum_{j=1}^k \delta^{-\frac{2(k-j+1)\alpha}{1+(k-j+2)\alpha}} \|\Delta_{x_k}^{\frac{\alpha}{2(1+\alpha)}} u\|_2^{\frac{2(1+\alpha)}{1+(k-j+2)\alpha}} \|f\|_2^{\frac{2(k-j+1)\alpha}{1+(k-j+2)\alpha}} \\
&\leq \frac{1}{4} \|\Delta_{x_{k+1}}^{\frac{\alpha}{2(1+\alpha)}} u\|_2^2 + C\delta^{-2} \|f\|_2^2.
\end{aligned}$$

Next we treat the trouble term $K_2(\delta)$. Let

$$\xi^{(k)} := (\xi_1, \xi_2, \dots, \xi_k), \quad \xi^{(k)} := (\xi_{k+1}, \xi_{k+2}, \dots, \xi_n)$$

and define

$$h_\delta(s, \xi^{(k)}) := \int_{\mathbb{R}^{kd}} (|\hat{u}|^2 \phi_\delta)(s, \xi, \xi^{(k)}) d\xi^{(k)}.$$

Integrating both sides of (3.4) with respect to variable $\xi^{(k)}$, we get

$$\begin{aligned}
&\partial_s h_\delta - A_{n-k} \xi^{(k)} \cdot \nabla_{\xi^{(k)}} h_\delta - \lambda h_\delta \\
&= \int_{\mathbb{R}^{kd}} \left(\sum_{j=1}^k \xi_j \nabla_{\xi_{j+1}} (|\hat{u}|^2 \phi_\delta) - g_{\phi_\delta} \right) d\xi^{(k)} =: g_{\phi_\delta}^{(k)},
\end{aligned}$$

and

$$h_\delta(s, \xi^{(k)}) = \int_0^\infty e^{-\lambda t} g_{\phi_\delta}^{(k)}(t + s, e^{-tA_{n-k}^*} \xi^{(k)}) dt.$$

Let

$$\psi_\varepsilon(\xi^{(k)}) := \chi(\varepsilon |\xi_2^{(k)}| / |\xi_1^{(k)}|^{1/(1+\alpha)}) = \chi(\varepsilon |\xi_{k+2}| / |\xi_{k+1}|^{1/(1+\alpha)}).$$

We make the following decomposition:

$$\begin{aligned}
K_2(\delta) &= \int_{-\infty}^\infty \int_{\mathbb{R}^{(n-k)d}} |\xi_{k+1}|^{\frac{2\alpha}{1+\alpha}} (1 - \psi_\varepsilon(\xi^{(k)})) h_\delta(s, \xi^{(k)}) d\xi^{(k)} ds \\
&\quad + \int_{-\infty}^\infty \int_{\mathbb{R}^{(n-k)d}} |\xi_{k+1}|^{\frac{2\alpha}{1+\alpha}} \psi_\varepsilon(\xi^{(k)}) h_\delta(s, \xi^{(k)}) d\xi^{(k)} ds \\
&= K_{21}(\delta, \varepsilon) + K_{22}(\delta, \varepsilon).
\end{aligned}$$

For $K_{21}(\delta, \varepsilon)$, we have

$$\begin{aligned}
K_{21}(\delta, \varepsilon) &\leq \int_{-\infty}^\infty \int_{\mathbb{R}^{(n-k)d}} |\xi_{k+1}|^{\frac{2\alpha}{1+\alpha}} \mathbf{1}_{\{\varepsilon |\xi_{k+2}| > |\xi_{k+1}|^{1/(1+\alpha)}\}} h_\delta(s, \xi^{(k)}) d\xi^{(k)} ds \\
&\leq \varepsilon^{2\alpha} \int_{-\infty}^\infty \int_{\mathbb{R}^{nd}} |\xi_{k+2}|^{2\alpha} |\hat{u}|^2(s, \xi) d\xi ds = \varepsilon^{2\alpha} \|\Delta_{x_{k+2}}^{\frac{\alpha}{2}} u\|_2^2.
\end{aligned}$$

For $K_{22}(\delta, \varepsilon)$, in view of $\xi_1^{(k)} = \xi_{k+1}$, we have

$$\begin{aligned}
K_{22}(\delta, \varepsilon) &= \int_{-\infty}^\infty \int_{\mathbb{R}^{(n-k)d}} |\xi_1^{(k)}|^{\frac{2\alpha}{1+\alpha}} \psi_\varepsilon(\xi^{(k)}) \int_0^\infty e^{-\lambda t} g_{\phi}^{(k)}(t + s, e^{-tA_{n-k}^*} \xi^{(k)}) dt d\xi^{(k)} ds \\
&= \int_{-\infty}^\infty \int_{\mathbb{R}^{(n-k)d}} \left(\int_0^\infty |(e^{tA_{n-k}^*} \xi^{(k)})_1|^{\frac{2\alpha}{1+\alpha}} \psi_\varepsilon(e^{tA_{n-k}^*} \xi^{(k)}) e^{-\lambda t} dt \right) g_{\phi}^{(k)}(s, \xi^{(k)}) d\xi^{(k)} ds.
\end{aligned}$$

Letting

$$\begin{aligned}\gamma_\varepsilon(\xi^{(k)}) &:= \int_0^\infty |(e^{tA_{n-k}^*} \xi^{(k)})_1|^{\frac{2\alpha}{1+\alpha}} \psi_\varepsilon(e^{tA_{n-k}^*} \xi^{(k)}) e^{-\lambda t} dt \\ &= |\xi_{k+1}|^{\frac{2\alpha}{1+\alpha}} \int_0^\infty \chi(\varepsilon|\xi_{k+2} + t\xi_{k+1}|/|\xi_{k+1}|^{1/(1+\alpha)}) e^{-\lambda t} dt,\end{aligned}$$

and recalling

$$g_{\phi_\delta}^{(k)} = \int_{\mathbb{R}^{kd}} g_{\phi_\delta}(\xi^{(k)}, \cdot) d\xi^{(k)} + \sum_{j=1}^k \int_{\mathbb{R}^{kd}} \xi_j \cdot \nabla_{\xi_{j+1}} (|\hat{u}|^2 \phi_\delta) d\xi^{(k)},$$

we may write

$$\begin{aligned}K_{22}(\delta, \varepsilon) &= \int_{-\infty}^\infty \int_{\mathbb{R}^{(n-k)d}} \gamma_\varepsilon(\xi^{(k)}) \int_{\mathbb{R}^d} g_{\phi_\delta}(s, \xi^{(k)}, \xi^{(k)}) d\xi^{(k)} d\xi^{(k)} ds \\ &\quad + \sum_{j=1}^k \int_{-\infty}^\infty \int_{\mathbb{R}^{(n-k)d}} \gamma_\varepsilon(\xi^{(k)}) \int_{\mathbb{R}^d} \xi_j \cdot \nabla_{\xi_{j+1}} (|\hat{u}|^2 \phi_\delta) d\xi^{(k)} d\xi^{(k)} ds \\ &=: K_{221}(\delta, \varepsilon) + K_{222}(\delta, \varepsilon).\end{aligned}$$

As in estimating (3.8), we have

$$\gamma_\varepsilon(\xi^{(k)}) \leq 4|\xi_{k+1}|^{\frac{\alpha}{1+\alpha}} / \varepsilon,$$

and also by the definition of g_{ϕ_δ}

$$|g_{\phi_\delta}| \leq 2|\hat{f}| |\hat{u}| + C\delta |\xi_{k+1}|^{\frac{\alpha}{1+\alpha}} |\hat{u}|^2.$$

Hence,

$$\begin{aligned}K_{221}(\delta, \varepsilon) &\leq \frac{8}{\varepsilon} \int_{-\infty}^\infty \int_{\mathbb{R}^{nd}} |\xi_{k+1}|^{\frac{\alpha}{1+\alpha}} |\hat{f}| |\hat{u}| + \frac{C\delta}{\varepsilon} \int_{-\infty}^\infty \int_{\mathbb{R}^{nd}} |\xi_{k+1}|^{\frac{2\alpha}{1+\alpha}} |\hat{u}|^2 \\ &\leq \left(\frac{1}{4} + \frac{C\delta}{\varepsilon}\right) \|\Delta_{x_{k+1}}^{\frac{\alpha}{2(1+\alpha)}} u\|_2^2 + C\varepsilon^{-2} \|f\|_2^2.\end{aligned}$$

Moreover, by elementary calculations, we also have

$$|\nabla_{\xi_{k+1}} \gamma_\varepsilon(\xi^{(k)})| \leq C|\xi_{k+1}|^{-\frac{1}{1+\alpha}} / \varepsilon.$$

Thus, since \hat{u} has compact support, by the integration by parts formula, we have

$$\begin{aligned}|K_{222}(\delta, \varepsilon)| &= \left| \int_{-\infty}^\infty \int_{\mathbb{R}^{(n-k)d}} \int_{\mathbb{R}^{kd}} \xi_k \nabla_{\xi_{k+1}} \gamma_\varepsilon(\xi^{(k)}) (|\hat{u}|^2 \phi_\delta) d\xi^{(k)} d\xi^{(k)} ds \right| \\ &\leq \frac{C}{\varepsilon} \int_{-\infty}^\infty \int_{\mathbb{R}^{nd}} |\xi_k| |\xi_{k+1}|^{-\frac{1}{1+\alpha}} (|\hat{u}|^2 \phi_\delta) d\xi ds \\ &\leq \frac{C}{\varepsilon} \int_{-\infty}^\infty \int_{\mathbb{R}^{nd}} |\xi_k| |\xi_{k+1}|^{-\frac{1}{1+\alpha}} |\hat{u}|^2 \mathbf{1}_{\{|\xi_k| \leq \delta |\xi_{k+1}|^{(1+2\alpha)/(1+\alpha)}\}} d\xi ds \\ &\leq \frac{C\delta}{\varepsilon} \int_{-\infty}^\infty \int_{\mathbb{R}^{nd}} |\xi_{k+1}|^{\frac{2\alpha}{1+\alpha}} |\hat{u}|^2 ds = \frac{C\delta}{\varepsilon} \|\Delta_{x_{k+1}}^{\frac{\alpha}{2(1+\alpha)}} u\|_2^2.\end{aligned}$$

Combining the above calculations, we obtain

$$\|\Delta_{x_{k+1}}^{\frac{\alpha}{2(1+\alpha)}} u\|_2^2 \leq \left(\frac{1}{4} + \frac{C'_1 \delta}{\varepsilon}\right) \|\Delta_{x_{k+1}}^{\frac{\alpha}{2(1+\alpha)}} u\|_2^2 + \varepsilon^{2\alpha} \|\Delta_{x_{k+2}}^{\frac{\alpha}{2}} u\|_2^2 + C'_2(\varepsilon^{-2} + \delta^{-2}) \|f\|_2^2.$$

Choosing $\delta = \varepsilon/(4C'_1)$, we get

$$\|\Delta_{x_{k+1}}^{\frac{\alpha}{2(1+\alpha)}} u\|_2^2 \leq 2\varepsilon^{2\alpha} \|\Delta_{x_{k+2}}^{\frac{\alpha}{2}} u\|_2^2 + 2C'_2(1 + 16(C'_1)^2)\varepsilon^{-2} \|f\|_2^2,$$

which then yields (3.6) for $j = k + 1$ by letting $\varepsilon = (\|f\|_2 / \|\Delta_{x_{k+2}}^{\frac{\alpha}{2}} u\|_2)^{\frac{1}{1+\alpha}}$. \square

4. PROOF OF THEOREM 1.1

4.1. **Case:** $p = 2$. Without loss of generality, we may assume $f \in C_c^\infty(\mathbb{R}^{1+nd})$. It follows from Fourier's transformation and Hölder's inequality that

$$\begin{aligned} & \int_{-\infty}^{\infty} \left\| \Delta_{x_n} \int_0^{\infty} e^{-\lambda t} \mathcal{T}_{s,t+s} f(t+s, \cdot) dt \right\|_2^2 ds \\ &= \int_{-\infty}^{\infty} \int_{\mathbb{R}^{nd}} |\xi_n|^2 \left| \int_0^{\infty} e^{-\lambda t} \widehat{\mathcal{T}_{s,t+s} f}(t+s, \xi) dt \right|^2 d\xi ds \\ &= \int_{-\infty}^{\infty} \int_{\mathbb{R}^{nd}} |\xi_n|^2 \left| \int_0^{\infty} e^{-\lambda t} e^{-\frac{1}{2} \int_0^t |(\sigma_{s+r}^a)^* e^{-rA^*} \xi|^2 dr} \widehat{f}(t+s, e^{-tA^*} \xi) dt \right|^2 d\xi ds \\ &\leq \int_{-\infty}^{\infty} \int_{\mathbb{R}^{nd}} \left(\int_0^{\infty} |\xi_n|^2 e^{-\frac{1}{2} \int_0^t |(\sigma_{s+r}^a)^* e^{-rA^*} \xi|^2 dr} |\widehat{f}(t+s, e^{-tA^*} \xi)|^2 dt \right) \\ &\quad \times \left(\int_0^{\infty} |\xi_n|^2 e^{-\frac{1}{2} \int_0^t |(\sigma_{s+r}^a)^* e^{-rA^*} \xi|^2 dr} dt \right) d\xi ds. \end{aligned}$$

Let Θ_t be the dilation operator defined by (1.7). By definitions (1.5), (1.6) and (1.7), it is easy to see that

$$t^{1/2} (\sigma_u^a)^* e^{-rA^*} \xi = (\sigma_u^a)^* e^{-rA^*} \Theta_{t^{1/2}}(\xi).$$

Thus, by the change of variables and as in showing (2.3), we have for some $c > 0$,

$$\int_0^t |(\sigma_{s+r}^a)^* e^{-rA^*} \xi|^2 dr = \int_0^1 |(\sigma_{s+tr}^a)^* e^{-rA^*} \Theta_{t^{1/2}}(\xi)|^2 dr \geq c |\Theta_{t^{1/2}} \xi|^2,$$

and similarly,

$$\int_0^t |(\sigma_{s+r}^a)^* e^{(t-r)A^*} \xi|^2 dr \geq c |\Theta_{t^{1/2}} \xi|^2.$$

Hence,

$$\int_0^{\infty} |\xi_n|^2 e^{-\frac{1}{2} \int_0^t |(\sigma_{s+r}^a)^* e^{-rA^*} \xi|^2 dr} dt \leq \int_0^{\infty} |\xi_n|^2 e^{-c |\Theta_{t^{1/2}} \xi|^2} dt \leq \int_0^{\infty} |\xi_n|^2 e^{-ct |\xi_n|^2} dt = c^{-1},$$

and

$$\begin{aligned} & \int_0^{\infty} |(e^{tA^*} \xi)_n|^2 e^{-\frac{1}{2} \int_0^t |(\sigma_{s-t+r}^a)^* e^{(t-r)A^*} \xi|^2 dr} dt \leq \int_0^{\infty} |(e^{tA^*} \xi)_n|^2 e^{-c |\Theta_{t^{1/2}} \xi|^2} dt \\ & \leq \sum_{j=1}^n \frac{1}{(n-j)!} \int_0^{\infty} |t^{n-j} \xi_j|^2 e^{-ct^{2(n-j)+1} |\xi_j|^2} dt \leq 2c^{-1}. \end{aligned}$$

Thus, by the change of variables and Fubini's theorem, we further have

$$\begin{aligned}
& \int_{-\infty}^{\infty} \left\| \Delta_{x_n} \int_0^{\infty} e^{-\lambda t} \mathcal{T}_{s,t+s} f(t+s, \cdot) dt \right\|_2^2 ds \\
& \leq c^{-1} \int_{-\infty}^{\infty} \int_{\mathbb{R}^{nd}} \left(\int_0^{\infty} |\xi_n|^2 e^{-\frac{1}{2} \int_0^t |(\sigma_{s+r}^a)^* e^{-rA^*} \xi|^2 dr} |\hat{f}(t+s, e^{-tA^*} \xi)|^2 dt \right) d\xi ds \\
& = c^{-1} \int_{-\infty}^{\infty} \int_{\mathbb{R}^{nd}} \left(\int_0^{\infty} |(e^{tA^*} \xi)_n|^2 e^{-\frac{1}{2} \int_0^t |(\sigma_{s+r}^a)^* e^{(t-r)A^*} \xi|^2 dr} |\hat{f}(t+s, \xi)|^2 dt \right) d\xi ds \\
& = c^{-1} \int_{-\infty}^{\infty} \int_{\mathbb{R}^{nd}} \left(\int_0^{\infty} |(e^{tA^*} \xi)_n|^2 e^{-\frac{1}{2} \int_0^t |(\sigma_{s+r}^a)^* e^{(t-r)A^*} \xi|^2 dr} dt \right) |\hat{f}(s, \xi)|^2 d\xi ds \\
& \leq 2c^{-2} \int_{-\infty}^{\infty} \int_{\mathbb{R}^{nd}} |\hat{f}(s, \xi)|^2 d\xi ds = 2c^{-2} \|f\|_2^2.
\end{aligned}$$

The proof of (1.10) for $p = 2$ is thus complete by (3.7).

4.2. **Case:** $p \in (2, \infty)$. For $j = 1, \dots, n$ and $\varepsilon \in (0, 1)$, let us define

$$\mathcal{P}_j^\varepsilon f := \mathcal{P}_j^a f_\varepsilon(s, x) := \Delta_{x_j}^{\frac{1}{1+2(n-j)}} \int_0^{\infty} e^{-\lambda t} \mathcal{T}_{s,t+s}^a f_\varepsilon(t+s, x) dt,$$

where $f_\varepsilon(t, x) = f(t) * \varrho_\varepsilon(x)$, $\varrho_\varepsilon(x) = \varepsilon^{-nd} \varrho(x/\varepsilon)$ is a family of mollifiers with $\varrho \in C_c^\infty(\mathbb{R}^{nd})$ and $\int \varrho = 1$, and the superscript a denotes the dependence on the diffusion coefficient a . To use Theorem 2.4, our main task is to show that $\mathcal{P}_j^\varepsilon$ is a bounded linear operator from $L^\infty(\mathbb{R}^{1+nd})$ to BMO . More precisely, we want to prove that for any $f \in L^\infty(\mathbb{R}^{1+nd})$ with $\|f\|_\infty \leq 1$, and any $Q = Q_r(t_0, x_0) \in \mathbb{Q}$,

$$\int_Q |\mathcal{P}_j^a f_\varepsilon(s, x) - c_j^Q|^2 \leq C, \tag{4.1}$$

where c_j^Q is a constant depending on Q and f_ε , and C only depends on n, κ, p, d , and not on ε .

Lemma 4.1. (*Scaling Property*) For any $Q = Q_r(t_0, x_0) \in \mathbb{Q}$, we have

$$\int_{Q_r(t_0, x_0)} |\mathcal{P}_j^a f_\varepsilon(s, x) - c|^2 = \int_{Q_1(0)} |\mathcal{P}_j^{\tilde{a}} \tilde{f}_\varepsilon(s, x) - c|^2, \tag{4.2}$$

where $c \in \mathbb{R}$, $\tilde{a}_s := a_{r^2 s + t_0}$ and $\tilde{f}_\varepsilon(t, x) := f_\varepsilon(r^2 t + t_0, \Theta_r x + e^{tA} x_0)$. Here Θ_r is the dilation operator defined in (1.7).

Proof. Let us write

$$u_\varepsilon(s, x) := \int_0^{\infty} e^{-\lambda t} \mathcal{T}_{s,t+s}^a f_\varepsilon(t+s, x) dt.$$

By the change of variables, we have

$$\tilde{u}_\varepsilon(s, x) := r^{-2} u_\varepsilon(r^2 s + t_0, \Theta_r x + e^{sA} x_0) = \int_0^{\infty} e^{-\lambda t} \mathcal{T}_{s,t+s}^{\tilde{a}} \tilde{f}_\varepsilon(t+s, x) dt.$$

Noticing that

$$(\Delta_{x_j}^{\frac{1}{1+2(n-j)}} \tilde{u}_\varepsilon)(s, x) = (\Delta_{x_j}^{\frac{1}{1+2(n-j)}} u_\varepsilon)(r^2 s + t_0, \Theta_r x + e^{sA} x_0),$$

by the change of variables again, we have

$$\int_{Q_r(t_0, x_0)} |\Delta_{x_j}^{\frac{1}{1+2(n-j)}} u_\varepsilon(s, x) - c|^2 = \int_{Q_1(0)} |\Delta_{x_j}^{\frac{1}{1+2(n-j)}} \tilde{u}_\varepsilon(s, x) - c|^2.$$

The proof is finished. \square

Noticing that

$$\int_0^\infty e^{-\lambda t} \mathcal{T}_{s,t+s}^a f_\varepsilon(t+s, x) dt = \int_s^\infty e^{\lambda(s-t)} \mathcal{T}_{s,t}^a f_\varepsilon(t, x) dt,$$

for $s \in [-1, 1]$, we split $\mathcal{P}_j^\varepsilon f(s, x) = \mathcal{P}_{j1}^\varepsilon f(s, x) + \mathcal{P}_{j2}^\varepsilon f(s, x)$ with

$$\begin{aligned} \mathcal{P}_{j1}^\varepsilon f(s, x) &:= \Delta_{x_j}^{\frac{1}{1+2(n-j)}} \int_s^2 e^{\lambda(s-t)} \mathcal{T}_{s,t} f_\varepsilon(t, x) dt, \\ \mathcal{P}_{j2}^\varepsilon f(s, x) &:= \Delta_{x_j}^{\frac{1}{1+2(n-j)}} \int_2^\infty e^{\lambda(s-t)} \mathcal{T}_{s,t} f_\varepsilon(t, x) dt. \end{aligned}$$

Convention: Without extra declaration, all the constants contained in “ \lesssim ” will depend only on n, κ, p, d .

Lemma 4.2. *Under (1.2), there is a constant $C = C(n, \kappa, p, d) > 0$ such that for all $f \in L^\infty(\mathbb{R}^{1+nd})$ with $\|f\|_\infty \leq 1$,*

$$\sup_{\varepsilon \in (0,1)} \int_{Q_1(0)} |\mathcal{P}_{j1}^\varepsilon f(s, x)|^2 \leq C. \quad (4.3)$$

Proof. For $s \in [-1, 1]$, let

$$u_\varepsilon(s, x) := \int_s^2 e^{\lambda(s-t)} \mathcal{T}_{s,t} f_\varepsilon(t, x) dt = \int_s^\infty e^{\lambda(s-t)} \mathcal{T}_{s,t} ((1_{[-1,2]} f_\varepsilon)(t))(x) dt.$$

Since $\|f\|_\infty \leq 1$, we have

$$\|u_\varepsilon(s)\|_\infty \leq 3, \quad s \in [-1, 1]. \quad (4.4)$$

By (2.6), we have for any $s \in [-1, 1]$,

$$\|\nabla_{x_n} u_\varepsilon(s)\|_\infty \leq \int_s^2 \|\nabla_{x_n} \mathcal{T}_{s,t} f_\varepsilon(t)\|_\infty dt \lesssim \int_s^2 (t-s)^{-1/2} dt \lesssim 1. \quad (4.5)$$

Let φ be a nonnegative smooth cutoff function in \mathbb{R}^{nd} with $\varphi(x) = 1$ for $|x| \leq n$ and $\varphi(x) = 0$ for $|x| > 2n$. Notice that

$$\partial_s u_\varepsilon + \mathcal{L}_s u_\varepsilon + f_\varepsilon 1_{[-1,2]}(s) = 0.$$

It is easy to see that $u_\varepsilon \varphi$ satisfies

$$\partial_s(u_\varepsilon \varphi) + \mathcal{L}_s(u_\varepsilon \varphi) = \mathcal{L}_s(u_\varepsilon \varphi) - (\mathcal{L}_s u_\varepsilon) \varphi - f_\varepsilon \varphi 1_{[-1,2]}(s) =: g_\varepsilon^\varphi,$$

which implies by (1.9) that

$$(u_\varepsilon \varphi)(s, x) = \int_s^\infty e^{\lambda(s-t)} \mathcal{T}_{s,t} g_\varepsilon^\varphi(t, x) dt.$$

By the definition of $\mathcal{P}_{j1}^\varepsilon$, we have

$$\begin{aligned} \int_{Q_1(0)} |\mathcal{P}_{j1}^\varepsilon f|^2 &= \int_{Q_1(0)} |\Delta_{x_j}^{\frac{1}{1+2(n-j)}} u_\varepsilon|^2 \leq \int_{\mathbb{R}^{1+nd}} |(\Delta_{x_j}^{\frac{1}{1+2(n-j)}} u_\varepsilon) \varphi|^2 \\ &\leq 2 \left\| \Delta_{x_j}^{\frac{1}{1+2(n-j)}} (u_\varepsilon \varphi) \right\|_2^2 + 2 \left\| \Delta_{x_j}^{\frac{1}{1+2(n-j)}} (u_\varepsilon \varphi) - (\Delta_{x_j}^{\frac{1}{1+2(n-j)}} u_\varepsilon) \varphi \right\|_2^2 =: I_1 + I_2. \end{aligned}$$

For I_1 , by (1.10) for $p = 2$ and (4.4), (4.5), we have

$$I_1 = \left\| \Delta_{x_j}^{\frac{1}{1+2(n-j)}} \int_s^\infty e^{\lambda(s-t)} \mathcal{T}_{s,t} g_\varepsilon^\varphi(t, x) dt \right\|_2 \lesssim \|g_\varepsilon^\varphi\|_2 \lesssim 1.$$

For I_2 , if $j = n$, then by (4.4) and (4.5),

$$I_2 = 2 \left\| u_\varepsilon \Delta_{x_n} \varphi + 2 \nabla_{x_n} u_\varepsilon \cdot \nabla_{x_n} \varphi \right\|_2^2 \lesssim 1;$$

if $j = 1, \dots, n-1$, then by definition (2.4) and (4.4),

$$\begin{aligned} I_2 &= \left\| \int_{\mathbb{R}^d} \frac{\delta_{z_j}^{(1)} u_\varepsilon \delta_{z_j}^{(1)} \varphi}{|z_j|^{d+2/(1+2(n-j))}} dz_j \right\|_2 \leq \int_{\mathbb{R}^d} \frac{\|\delta_{z_j}^{(1)} u_\varepsilon \delta_{z_j}^{(1)} \varphi\|_2}{|z_j|^{d+2/(1+2(n-j))}} dz_j \\ &\leq 6 \int_{\mathbb{R}^d} \frac{\|\delta_{z_j}^{(1)} \varphi\|_2}{|z_j|^{d+2/(1+2(n-j))}} dz_j \lesssim \int_{\mathbb{R}^d} \frac{1 \wedge |z_j|}{|z_j|^{d+2/(1+2(n-j))}} dz_j \lesssim 1. \end{aligned}$$

The proof is complete. \square

The next lemma is crucial for treating $\mathcal{P}_{j2}^\varepsilon f$.

Lemma 4.3. *Under (1.3), there is a constant $C = C(n, \kappa, p, d) > 0$ such that for all $f \in L^\infty(\mathbb{R}^{1+nd})$ with $\|f\|_\infty \leq 1$ and all $s \in [-1, 1]$,*

$$\sup_{\varepsilon \in (0,1)} \int_2^\infty \left| \Delta_{x_j}^{\frac{1}{1+2(n-j)}} \mathcal{T}_{s,t} f_\varepsilon(t, 0) - \Delta_{x_j}^{\frac{1}{1+2(n-j)}} \mathcal{T}_{0,t} f_\varepsilon(t, 0) \right| dt \leq C. \quad (4.6)$$

Proof. Noticing that by (1.9), for each $s \in (0, t)$ and $x \in \mathbb{R}^{nd}$,

$$\mathcal{T}_{0,t} f_\varepsilon(t, x) - \mathcal{T}_{s,t} f_\varepsilon(t, x) = \int_0^s \mathcal{L}_r \mathcal{T}_{r,t} f_\varepsilon(t, x) dr, \quad (4.7)$$

we have

$$\begin{aligned} I_j^\varepsilon(s, t) &:= \Delta_{x_j}^{\frac{1}{1+2(n-j)}} \mathcal{T}_{0,t} f_\varepsilon(t, 0) - \Delta_{x_j}^{\frac{1}{1+2(n-j)}} \mathcal{T}_{s,t} f_\varepsilon(t, 0) \\ &= \int_0^s (\Delta_{x_j}^{\frac{1}{1+2(n-j)}} \mathcal{L}_r \mathcal{T}_{r,t} f_\varepsilon)(t, 0) dr. \end{aligned}$$

For $j = 1$, we have for all $s \in [-1, 1]$,

$$\int_2^\infty |I_1^\varepsilon(s, t)| dt = \int_2^\infty \left| \int_0^s (\Delta_{x_1}^{\frac{1}{1+2(n-1)}} \text{tr}(a_r \cdot \nabla_{x_n}^2) \mathcal{T}_{r,t} f_\varepsilon)(t, 0) dr \right| dt$$

$$\begin{aligned}
&\leq \kappa \int_2^\infty \int_0^s |\Delta_{x_1}^{\frac{1}{1+2(n-1)}} \nabla_{x_n}^2 \mathcal{T}_{r,t} f_\varepsilon|(t, 0) dr dt \\
&\stackrel{(2.6)}{\lesssim} \int_2^\infty \int_0^s (t-r)^{-3/2} dr dt \lesssim 1.
\end{aligned}$$

For $j = n$, we have for all $s \in [-1, 1]$,

$$\begin{aligned}
\int_2^\infty |I_n^\varepsilon(s, t)| dt &= \int_2^\infty \left| \int_0^s ((\Delta_{x_n} \operatorname{tr}(a_r \cdot \nabla_{x_n}^2) + (\nabla_{x_n} \cdot \nabla_{x_{n-1}})) \mathcal{T}_{r,t} f_\varepsilon)(t, 0) dr \right| dt \\
&\leq \int_2^\infty \int_0^s (\kappa |\Delta_{x_n} \nabla_{x_n}^2 \mathcal{T}_{r,t} f_\varepsilon| + |(\nabla_{x_n} \cdot \nabla_{x_{n-1}}) \mathcal{T}_{r,t} f_\varepsilon|)(t, 0) dr dt \\
&\stackrel{(2.6)}{\lesssim} \int_2^\infty \int_0^s (t-r)^{-2} dr dt \lesssim 1.
\end{aligned}$$

For $j = 2, \dots, n-1$, since $\Delta_{x_j}^{\frac{1}{1+2(n-j)}}$ is a nonlocal operator, we have to carefully treat the trouble term $\Delta_{x_j}^{\frac{1}{1+2(n-j)}}(x_j \cdot \nabla_{x_{j-1}} \mathcal{T}_{r,t} f_\varepsilon)(t, 0)$. Fix

$$\gamma \in \left(\frac{2(n-j)+1}{2}, \frac{(2(n-j)+1)^2}{4(n-j)} \right).$$

By (2.4) and (4.7), we may write

$$\begin{aligned}
I_j^\varepsilon(s, t) &= \int_{|z_j| > t^\gamma} (\delta_{z_j} \mathcal{T}_{0,t} f_\varepsilon(t, 0) - \delta_{z_j} \mathcal{T}_{s,t} f_\varepsilon(t, 0)) \frac{dz_j}{|z_j|^{d+2/(1+2(n-j))}} \\
&\quad + \int_{|z_j| \leq t^\gamma} \left(\int_0^s \delta_{z_j} \operatorname{tr}(a_r \cdot \nabla_{x_n}^2) \mathcal{T}_{r,t} f_\varepsilon(t, 0) dr \right) \frac{dz_j}{|z_j|^{d+2/(1+2(n-j))}} \\
&\quad + \int_{|z_j| \leq t^\gamma} \left(\int_0^s \delta_{z_j} (Ax \cdot \nabla \mathcal{T}_{r,t} f_\varepsilon)(t, 0) dr \right) \frac{dz_j}{|z_j|^{d+2/(1+2(n-j))}} \\
&=: I_{j1}^\varepsilon(s, t) + I_{j2}^\varepsilon(s, t) + I_{j3}^\varepsilon(s, t).
\end{aligned}$$

For $I_{j1}^\varepsilon(s, t)$, thanks to $\gamma > \frac{2(n-j)+1}{2}$, we have for all $s \in [-1, 1]$,

$$\int_2^\infty |I_{j1}^\varepsilon(s, t)| dt \lesssim \int_2^\infty \int_{|z_j| > t^\gamma} \frac{dz_j}{|z_j|^{d+2/(1+2(n-j))}} dt \lesssim \int_2^\infty |t|^{-2\gamma/(1+2(n-j))} dt \lesssim 1.$$

For $I_{j2}^\varepsilon(s, t)$, by (2.6) and $\gamma < \frac{(2(n-j)+1)^2}{2(2(n-j)-1)}$, we have for all $s \in [-1, 1]$,

$$\begin{aligned}
\int_2^\infty |I_{j2}^\varepsilon(s, t)| dt &\lesssim \int_2^\infty \int_0^s \left(\int_{|z_j| \leq t^\gamma} |z_j| \|\nabla_{x_j} \nabla_{x_n}^2 \mathcal{T}_{r,t} f_\varepsilon\|_\infty \frac{dz_j}{|z_j|^{d+2/(1+2(n-j))}} \right) dr dt \\
&\lesssim \int_2^\infty \int_0^s (t-r)^{-(2(n-j)+3)/2} \left(\int_{|z_j| \leq t^\gamma} \frac{|z_j| dz_j}{|z_j|^{d+2/(1+2(n-j))}} \right) dr dt \\
&\lesssim \int_2^\infty (t-1)^{-(2(n-j)+3)/2} t^{\gamma(1-2/(1+2(n-j)))} dt \lesssim 1.
\end{aligned}$$

For $I_{j3}^\varepsilon(s, t)$, letting $\tilde{z}_j = (0, \dots, 0, z_j, 0, \dots, 0)$ and observing that

$$\int_{|z_j| \leq \rho'} \frac{z_j dz_j}{|z_j|^{d+2/(1+2(n-j))}} = 0,$$

by (2.6) and $\gamma < \frac{(2(n-j)+1)^2}{4(n-j)}$, we have for all $s \in [-1, 1]$,

$$\begin{aligned} \int_2^\infty |I_{j3}^\varepsilon(s, t)| dt &= \int_2^\infty \left| \int_0^s \int_{|z_j| \leq \rho'} z_j \cdot \nabla_{x_{j-1}} \mathcal{T}_{r,t} f_\varepsilon(t, \tilde{z}_j) \frac{dz_j}{|z_j|^{d+2/(1+2(n-j))}} dr \right| dt \\ &= \int_2^\infty \left| \int_0^s \int_{|z_j| \leq \rho'} z_j \cdot \nabla_{x_{j-1}} (\mathcal{T}_{r,t} f_\varepsilon(t, \tilde{z}_j) - \mathcal{T}_{r,t} f_\varepsilon(t, 0)) \frac{dz_j}{|z_j|^{d+2/(1+2(n-j))}} dr \right| dt \\ &\leq \int_2^\infty \left| \int_0^s \int_{|z_j| \leq \rho'} |z_j|^2 \|\nabla_{x_j} \nabla_{x_{j-1}} \mathcal{T}_{r,t} f_\varepsilon\|_\infty \frac{dz_j}{|z_j|^{d+2/(1+2(n-j))}} dr \right| dt \\ &\lesssim \int_2^\infty \left(\int_0^s (t-r)^{-2(n-j)-2} \int_{|z| \leq \rho'} \frac{|z_j|^2 dz_j}{|z_j|^{d+2/(1+2(n-j))}} dr \right) dt \\ &\lesssim \int_2^\infty (t-1)^{-2(n-j)-2} \rho'^{(2-2/(1+2(n-j)))} dt \lesssim 1. \end{aligned}$$

Combining the above calculations, we obtain the desired estimate. \square

Lemma 4.4. *Under (1.3), there is a constant $C > 0$ depending only on n, κ, p, d such that for all $f \in L^\infty(\mathbb{R}^{1+nd})$ with $\|f\|_\infty \leq 1$,*

$$\sup_{\varepsilon \in (0,1)} \int_{Q_1(0)} |\mathcal{P}_{j2}^\varepsilon f(s, x) - \mathcal{P}_{j2}^\varepsilon f(0, 0)|^2 \leq C. \quad (4.8)$$

Proof. By definition, we have

$$\begin{aligned} |\mathcal{P}_{j2}^\varepsilon f(s, x) - \mathcal{P}_{j2}^\varepsilon f(0, 0)| &\leq \int_2^\infty |e^{\lambda(s-t)} - e^{-\lambda t}| \|\Delta_{x_j}^{\frac{1}{1+2(n-j)}} \mathcal{T}_{s,t} f_\varepsilon(t)\|_\infty dt \\ &+ \int_2^\infty e^{-\lambda t} |\Delta_{x_j}^{\frac{1}{1+2(n-j)}} \mathcal{T}_{s,t} f_\varepsilon(t, x) - \Delta_{x_j}^{\frac{1}{1+2(n-j)}} \mathcal{T}_{s,t} f_\varepsilon(t, 0)| dt \\ &+ \int_2^\infty e^{-\lambda t} |\Delta_{x_j}^{\frac{1}{1+2(n-j)}} \mathcal{T}_{s,t} f_\varepsilon(t, 0) - \Delta_{x_j}^{\frac{1}{1+2(n-j)}} \mathcal{T}_{0,t} f_\varepsilon(t, 0)| dt \\ &=: I_1(s) + I_2(s, x) + I_3(s). \end{aligned}$$

Noticing that by (2.6),

$$\|\Delta_{x_j}^{\frac{1}{1+2(n-j)}} \mathcal{T}_{s,t} f_\varepsilon(t)\|_\infty \lesssim (t-s)^{-1},$$

$$\|\nabla_{x_k} \Delta_{x_j}^{\frac{1}{1+2(n-j)}} \mathcal{T}_{s,t} f_\varepsilon(t)\|_\infty \lesssim (t-s)^{-(2(n-k)+3)/2},$$

we have for all $s \in [-1, 1]$,

$$\begin{aligned} I_1 &\lesssim \int_2^\infty |e^{\lambda(s-t)} - e^{-\lambda t}| (t-s)^{-1} dt \\ &\lesssim |e^{\lambda s} - 1| \int_2^\infty e^{-\lambda t} dt = |e^{\lambda s} - 1| e^{-2\lambda} / \lambda \lesssim 1, \end{aligned}$$

and for all $(s, x) \in Q_1(0)$,

$$I_2(s, x) \lesssim \int_2^\infty \sum_{k=1}^n (t-s)^{-(2(n-k)+3)/2} dt \lesssim 1.$$

Moreover, by (4.6), we have for all $s \in [-1, 1]$,

$$I_3(s) \lesssim 1.$$

Combining the above calculations, we obtain (4.8). \square

Now we can give

Proof of (1.10) for $p \in (2, \infty)$. By Lemmas 4.1, 4.2 and 4.4, it is easy to see that $\mathcal{P}_j^\varepsilon : L^\infty(\mathbb{R}^{1+nd}) \rightarrow BMO$ is a bounded linear operator with bound independent of ε . Estimate (1.10) for $p \in (2, \infty)$ follows by Theorem 2.4 and the well-proved estimate for $p = 2$. \square

4.3. Case: $p \in (1, 2)$. We shall use duality argument to show (1.10) for $p \in (1, 2)$. Let $\mathcal{T}_{s,t}^*$ be the adjoint operator of $\mathcal{T}_{s,t}$, that is,

$$\int g \mathcal{T}_{s,t} f = \int f \mathcal{T}_{s,t}^* g.$$

By the definition of $\mathcal{T}_{s,t}$, we have

$$\mathcal{T}_{s,t}^* f(x) := \mathcal{T}_{s,t}^{*,a} f(x) := \mathbb{E} f \left(e^{(s-t)A} x - \int_s^t e^{(s-r)A} \sigma_r^a dW_r \right).$$

For $j = 1, \dots, n$, we introduce a new operator

$$\mathcal{Q}_j^\varepsilon f := \mathcal{Q}_j^a f_\varepsilon(s, x) := \int_{-\infty}^t e^{\lambda(s-t)} \mathcal{T}_{s,t}^{*,a} \Delta_{x_j}^{\frac{1}{1+2(n-j)}} f_\varepsilon(s, x) ds,$$

where $f_\varepsilon(t, x) = f(t, \cdot) * \varrho_\varepsilon(x)$ so that $\mathcal{Q}_j^\varepsilon f$ is well defined for $f \in L^\infty(\mathbb{R}^{1+nd})$. Notice that \mathcal{Q}_j^a can be considered as the adjoint operator of \mathcal{P}_j^a in the sense that

$$\int \mathcal{P}_j^a f g = \int \mathcal{Q}_j^a g f.$$

As in the previous subsection, we want to show that

$$\mathcal{Q}_j^\varepsilon \text{ is a bounded linear operator from } L^\infty(\mathbb{R}^{1+nd}) \text{ to } BMO.$$

First of all, as in Lemma 4.1 we have

$$\int_{Q_r(t_0, x_0)} |\mathcal{Q}_j^a f_\varepsilon(s, x) - c|^2 = \int_{Q_1(0)} |\mathcal{Q}_j^{\tilde{a}} \tilde{f}_\varepsilon(s, x) - c|^2,$$

where \tilde{a} and \tilde{f} are defined as in Lemma 4.1. We aim to prove that there is a constant $C = C(n, \kappa, p, d) > 0$ independent of $\varepsilon \in (0, 1)$ such that for all $f \in L^\infty(\mathbb{R}^{1+nd})$ with $\|f\|_\infty \leq 1$,

$$\int_{Q_1(0)} |\mathcal{Q}_j^{\tilde{a}} \tilde{f}_\varepsilon(s, x) - c|^2 \leq C.$$

Below we drop \tilde{a} and the tilde, and make the following decomposition as above,

$$\mathcal{Q}_j^\varepsilon f(t, x) = \left(\int_{-2}^t + \int_{-\infty}^{-2} \right) e^{\lambda(s-t)} \mathcal{T}_{s,t}^* \Delta_{x_j}^{\frac{1}{1+2(n-j)}} f_\varepsilon(s, x) ds =: \mathcal{Q}_{j1}^\varepsilon f(t, x) + \mathcal{Q}_{j2}^\varepsilon f(t, x).$$

The following lemma is crucial for treating $\mathcal{Q}_{j1}^\varepsilon$.

Lemma 4.5. *Let $\varphi \in C_c^\infty(\mathbb{R}^{nd})$. For any $p \in [1, 2]$, there are constants $C_\varphi, \beta > 0$ such that for all $h \in L^2(\mathbb{R}^{nd})$ and $0 < t - s \leq 3$,*

$$\|\Delta_{x_j}^{\frac{1}{1+2(n-j)}} (\mathcal{T}_{s,t}(\varphi^2 h) - \varphi_{s,t} \mathcal{T}_{s,t}(\varphi h))\|_p \leq C_\varphi (t-s)^{\beta-1} \|h\|_2, \quad (4.9)$$

$$\|\Delta_{x_j}^{\frac{1}{1+2(n-j)}} (\varphi_{s,t} \mathcal{T}_{s,t}(\varphi h)) - \varphi_{s,t} \Delta_{x_j}^{\frac{1}{1+2(n-j)}} \mathcal{T}_{s,t}(\varphi h)\|_p \leq C_\varphi (t-s)^{\beta-1} \|h\|_2, \quad (4.10)$$

where $\varphi_{s,t}(x) := \varphi(e^{(t-s)A}x)$.

Proof. (1) Let $p_{s,t}^{(0)}(y)$ be the distribution density of $X_{s,t}(0)$. By the definition and the chain rule, we have

$$\begin{aligned} \nabla_{x_j} \mathcal{T}_{s,t} f(x) &= \int_{\mathbb{R}^{nd}} f(y) \nabla_{x_j} p_{s,t}^{(0)}(y - e^{(t-s)A} \cdot)(x) dy \\ &= \int_{\mathbb{R}^{nd}} f(y) \sum_{i=1}^j (\nabla_{y_i} p_{s,t}^{(0)})(y - e^{(t-s)A}x) \frac{(t-s)^{j-i}}{(j-i)!} dy \\ &= \int_{\mathbb{R}^{nd}} f(y + e^{(t-s)A}x) \sum_{i=1}^j (\nabla_{y_i} p_{s,t}^{(0)})(y) \frac{(t-s)^{j-i}}{(j-i)!} dy. \end{aligned}$$

By this formula, we have

$$\begin{aligned} &\|\nabla_{x_j} \mathcal{T}_{s,t}(\varphi^2 h) - \varphi_{s,t} \nabla_{x_j} \mathcal{T}_{s,t}(\varphi h)\|_p \\ &\leq \|\nabla \varphi\|_\infty \|\varphi h\|_p \sum_{i=1}^j \int_{\mathbb{R}^{nd}} |y| |\nabla_{y_i} p_{s,t}^{(0)}(y)| \frac{(t-s)^{j-i}}{(j-i)!} dy. \end{aligned} \quad (4.11)$$

By (2.1), we have

$$\begin{aligned} \int_{\mathbb{R}^{nd}} |y| |\nabla_{y_i} p_{s,t}^{(0)}(y)| dy &\lesssim (t-s)^{-(n^2 d + 2(n-i) + 1)/2} \sum_{k=1}^n \int_{\mathbb{R}^{nd}} |y_k| e^{-c|\Theta_{(t-s)^{-1/2}y}|^2} dy \\ &\lesssim \sum_{k=1}^n (t-s)^{-(n^2 d + 2(k-i))/2} \int_{\mathbb{R}^{nd}} e^{-c|\Theta_{(t-s)^{-1/2}y}|^2/2} dy \lesssim \sum_{k=1}^n (t-s)^{i-k}. \end{aligned}$$

Substituting this into (4.11), we obtain

$$\|\nabla_{x_j} \mathcal{T}_{s,t}(\varphi^2 h) - \varphi_{s,t} \nabla_{x_j} \mathcal{T}_{s,t}(\varphi h)\|_p \lesssim \|\varphi h\|_p (t-s)^{j-n} \lesssim \|h\|_2 (t-s)^{j-n},$$

and further,

$$\|\nabla_{x_j} (\mathcal{T}_{s,t}(\varphi^2 h) - \varphi_{s,t} \mathcal{T}_{s,t}(\varphi h))\|_p \lesssim \|h\|_2 (t-s)^{j-n}.$$

Hence, for $j = 1, 2, \dots, n-1$,

$$\|\Delta_{x_j}^{\frac{1}{1+2(n-j)}} (\mathcal{T}_{s,t}(\varphi^2 h) - \varphi_{s,t} \mathcal{T}_{s,t}(\varphi h))\|_p$$

$$\begin{aligned}
&\leq \int_{\mathbb{R}^d} \|\delta_{z_j}(\mathcal{T}_{s,t}(\varphi^2 h) - \varphi_{s,t} \mathcal{T}_{s,t}(\varphi h))\|_p \frac{dz_j}{|z_j|^{d+2/(1+2(n-j))}} \\
&\leq 2 \int_{|z_j| > (t-s)^{n-j}} (\|\mathcal{T}_{s,t}(\varphi^2 h)\|_p + \|\varphi_{s,t} \mathcal{T}_{s,t}(\varphi h)\|_p) \frac{dz_j}{|z_j|^{d+2/(1+2(n-j))}} \\
&\quad + \int_{|z_j| \leq (t-s)^{n-j}} (\|\nabla_{x_j}(\mathcal{T}_{s,t}(\varphi^2 h) - \varphi_{s,t} \mathcal{T}_{s,t}(\varphi h))\|_p) \frac{|z_j| dz_j}{|z_j|^{d+2/(1+2(n-j))}} \\
&\lesssim (\|\varphi^2 h\|_p + \|\varphi h\|_p) \int_{|z_j| > (t-s)^{n-j}} \frac{dz_j}{|z_j|^{d+2/(1+2(n-j))}} \\
&\quad + \|h\|_2 (t-s)^{j-n} \int_{|z_j| \leq (t-s)^{n-j}} \frac{|z_j| dz_j}{|z_j|^{d+2/(1+2(n-j))}} \lesssim \|h\|_2 (t-s)^{-\frac{2(n-j)}{1+2(n-j)}}.
\end{aligned}$$

Thus we get (4.9). For $j = n$, (4.9) is direct by the chain rule.

(2) For $j = 1, \dots, n-1$, by (2.7) we have

$$\begin{aligned}
&\|\Delta_{x_j}^{\frac{1}{1+2(n-j)}}(\varphi_{s,t} \mathcal{T}_{s,t}(\varphi h)) - \varphi_{s,t} \Delta_{x_j}^{\frac{1}{1+2(n-j)}} \mathcal{T}_{s,t}(\varphi h)\|_p \\
&\leq 2 \int_{\mathbb{R}^d} \frac{\|\delta_{z_j}^{(1)} \varphi_{s,t}\|_\infty \|\delta_{z_j}^{(1)} \mathcal{T}_{s,t}(\varphi h)\|_p}{|z_j|^{d+2/(1+2(n-j))}} dz_j \\
&\lesssim \|\varphi h\|_p \int_{\mathbb{R}^d} \frac{1 \wedge ((t-s)^{n-j} |z_j|)}{|z_j|^{d+2/(1+2(n-j))}} dz_j \lesssim \|h\|_2 (t-s)^{\frac{2(n-j)}{1+2(n-j)}},
\end{aligned}$$

which gives (4.10). For $j = n$, (4.10) is direct by the chain rule. \square

Lemma 4.6. *Under (1.3), there is a constant $C = C(n, \kappa, p, d) > 0$ such that for all $f \in L^\infty(\mathbb{R}^{1+nd})$ with $\|f\|_\infty \leq 1$,*

$$\sup_{\varepsilon \in (0,1)} \int_{Q_1(0)} |\mathcal{D}_{j_1}^\varepsilon f(s, x)|^2 \leq C. \quad (4.12)$$

Proof. For $t \in [-2, 1]$, define

$$u_\varepsilon(t, x) := \mathcal{D}_{j_1}^\varepsilon f(t, x) = \int_{-\infty}^t e^{\lambda(s-t)} \mathcal{T}_{s,t}^* \Delta_{x_j}^{\frac{1}{1+2(n-j)}} ((1_{[-2,1]} f_\varepsilon)(s))(x) ds.$$

Let φ be a nonnegative smooth cutoff function in \mathbb{R}^{nd} with $\varphi(x) = 1$ for $|x| \leq 1$ and $\varphi(x) = 0$ for $|x| > 2$. We have

$$\begin{aligned}
\|u\|_{L^2(Q_1(0))} &\leq \|1_{[-2,1]} u \varphi^2\|_2 = \sup_{\|h\|_2 \leq 1} \int_{\mathbb{R}^{1+nd}} 1_{[-2,1]} u \varphi^2 h \\
&= \sup_{\|h\|_2 \leq 1} \int_{\mathbb{R}^{1+nd}} 1_{[-2,1]} f_\varepsilon \Delta_{x_j}^{\frac{1}{1+2(n-j)}} \int_0^1 e^{\lambda(-t)} \mathcal{T}_{\cdot,t}(\varphi^2 h(t)) dt \\
&\leq \sup_{\|h\|_2 \leq 1} \left\| \int_{[-2,1]} \Delta_{x_j}^{\frac{1}{1+2(n-j)}} \int_0^1 e^{\lambda(s-t)} \mathcal{T}_{s,t}(\varphi^2 h(t)) dt \right\|_1.
\end{aligned}$$

Since the support of $\varphi_{s,t}(x) = \varphi(e^{(t-s)A}x)$ is contained in $\{x : |x| \leq n\}$ for $|t - s| \leq 3$, by (4.9) and (4.10), we have for any $h \in L^2(\mathbb{R}^{1+nd})$ with $\|h\|_2 \leq 1$.

$$\begin{aligned}
& \left\| 1_{[-2,1]} \Delta_{x_j}^{\frac{1}{1+2(n-j)}} \int_{\cdot}^1 e^{\lambda(s-t)} \mathcal{T}_{s,t}(\varphi^2 h(t)) dt \right\|_1 \\
& \stackrel{(4.9)}{\lesssim} \left\| 1_{[-2,1]} \Delta_{x_j}^{\frac{1}{1+2(n-j)}} \int_{\cdot}^1 e^{\lambda(s-t)} \varphi_{s,t} \mathcal{T}_{s,t}(\varphi h(t)) dt \right\|_1 + 1 \\
& \stackrel{(4.10)}{\lesssim} \left\| 1_{[-2,1]} \int_{\cdot}^1 e^{\lambda(s-t)} \varphi_{s,t} \Delta_{x_j}^{\frac{1}{1+2(n-j)}} \mathcal{T}_{s,t}(\varphi h(t)) dt \right\|_1 + 1 \\
& \lesssim \left\| 1_{[-2,1]} \int_{\cdot}^1 e^{\lambda(s-t)} \varphi_{s,t} \Delta_{x_j}^{\frac{1}{1+2(n-j)}} \mathcal{T}_{s,t}(\varphi h(t)) dt \right\|_2 + 1 \\
& \stackrel{(4.10)}{\lesssim} \left\| 1_{[-2,1]} \Delta_{x_j}^{\frac{1}{1+2(n-j)}} \int_{\cdot}^1 e^{\lambda(s-t)} \varphi_{s,t} \mathcal{T}_{s,t}(\varphi h(t)) dt \right\|_2 + 1 \\
& \stackrel{(4.9)}{\lesssim} \left\| \Delta_{x_j}^{\frac{1}{1+2(n-j)}} \int_{\cdot}^{\infty} e^{\lambda(s-t)} \mathcal{T}_{s,t}(1_{[-2,1]} \varphi^2 h(t)) dt \right\|_2 + 1 \\
& \lesssim \|\varphi^2 h\|_2 + 1 \lesssim 1,
\end{aligned}$$

where the last step is due to (1.10) for $p = 2$. The proof is complete. \square

The following lemma is the same as in Lemma 4.4.

Lemma 4.7. *Under (1.3), there is a constant $C = C(n, \kappa, p, d) > 0$ such that for all $f \in L^\infty(\mathbb{R}^{1+nd})$ with $\|f\|_\infty \leq 1$,*

$$\sup_{\varepsilon \in (0,1)} \int_{Q_1(0)} |\mathcal{Q}_{j_2}^\varepsilon f(t, x) - \mathcal{Q}_{j_2}^\varepsilon f(0, 0)|^2 \leq C.$$

Proof. By (2.7), we have for all $t \in [-1, 1]$,

$$\begin{aligned}
& \int_{-\infty}^{-2} |\mathcal{T}_{s,t}^* \Delta_{x_j}^{\frac{1}{1+2(n-j)}} f_\varepsilon(s, 0) - \mathcal{T}_{s,0}^* \Delta_{x_j}^{\frac{1}{1+2(n-j)}} f_\varepsilon(s, 0)| ds \\
& \leq \int_{-\infty}^{-2} \int_0^t |\partial_r \mathcal{T}_{s,r}^* \Delta_{x_j}^{\frac{1}{1+2(n-j)}} f_\varepsilon(s, 0)| dr ds \\
& = \int_{-\infty}^{-2} \int_0^t |\mathcal{L}_r^* \mathcal{T}_{s,r}^* \Delta_{x_j}^{\frac{1}{1+2(n-j)}} f_\varepsilon(s, 0)| dr ds \\
& = \int_{-\infty}^{-2} \int_0^t |\text{tr}(a_r \cdot \nabla_{x_n}^2) \mathcal{T}_{s,r}^* \Delta_{x_j}^{\frac{1}{1+2(n-j)}} f_\varepsilon(s, 0)| dr ds \\
& \leq C \int_{-\infty}^{-2} \int_0^t (r-s)^{-2} dr ds \leq C.
\end{aligned}$$

Using this estimate and (2.7), as in the proof of Lemma 4.4, we obtain the desired estimate. \square

Proof of (1.10) for $p \in (1, 2)$. By Lemmas 4.6 and 4.7, we know that

$$\mathcal{Q}_j^\varepsilon : L^\infty(\mathbb{R}^{1+2d}) \rightarrow BMO \text{ is bounded with norm independent of } \varepsilon.$$

Moreover, by duality, we also have

$$\mathcal{Q}_j^\varepsilon : L^2(\mathbb{R}^{1+2d}) \rightarrow L^2(\mathbb{R}^{1+2d}) \text{ is bounded with norm independent of } \varepsilon.$$

Hence, for $q = p/(p-1) \in (2, \infty)$, by Theorem 2.4, we have for some $C > 0$ independent of ε ,

$$\|\mathcal{Q}_j^\varepsilon f\|_q = \left\| \int_{-\infty}^t e^{\lambda(s-t)} \mathcal{T}_{s,t}^* \Delta_{x_j}^{\frac{1}{1+2(n-j)}} f_\varepsilon ds \right\|_q \leq C \|f\|_q.$$

Now for $p \in (1, 2)$, by Fatou's lemma, we get

$$\begin{aligned} \|\mathcal{P}_j f\|_p &\leq \|f\|_p \sup_{\|h\|_q \leq 1} \left\| \int_{-\infty}^t e^{\lambda(s-t)} \mathcal{T}_{s,t}^* \Delta_{x_j}^{\frac{1}{1+2(n-j)}} h ds \right\|_q \\ &\leq \|f\|_p \sup_{\|h\|_q \leq 1} \lim_{\varepsilon \rightarrow 0} \left\| \int_{-\infty}^t e^{\lambda(s-t)} \mathcal{T}_{s,t}^* \Delta_{x_j}^{\frac{1}{1+2(n-j)}} h_\varepsilon ds \right\|_q \leq C \|f\|_p, \end{aligned}$$

which gives (1.10) for $p \in (1, 2)$. □

REFERENCES

- [1] R. Alexander: Fractional order kinetic equations and hypoellipticity. *Anal. Appl.* (Singap.) **10**, no.3 (2012), 237-247.
- [2] F. Bouchut: Hypoelliptic regularity in kinetic equations. *J. Math. Pures Appl.* **81** (2002), 1135-1159.
- [3] M. Bramanti, G. Cupini, E. Lanconelli and E. Priola: Global L^p -estimate for degenerate Ornstein-Uhlenbeck operators. *Math Z.* **266** (2010), 789-816.
- [4] Z.-Q. Chen and X. Zhang: L^p -maximal hypoelliptic regularity of nonlocal kinetic Fokker-Planck operators. arXiv:1608.05502.
- [5] Fedrizzi E., Flandoli F., Priola E. and Vovelle J.: Regularity of Stochastic Kinetic Equations. <http://arxiv.org/pdf/1606.01088v2.pdf>.
- [6] Huang L., Menozzi S. and Priola E.: L^p -estimates for degenerate non-local Kolmogorov operators. arXiv:1607.08718v2.
- [7] A. N. Kolmogorov: Zufällige Bewegungen. *Ann. of Math.* **35** (1934), 116-117.
- [8] E. M. Stein: *Harmonic Analysis: Real-Variable Methods, Orthogonality and Oscillatory Integrals*. Princeton University Press, 1993.
- [9] X. Zhang: L^p -maximal regularity of nonlocal parabolic equations and applications. *Ann. I. H. Poincaré-AN* **30** (2013), 573-614.
- [10] X. Zhang: Stochastic Hamiltonian flows with singular coefficients. arXiv:1606.04360.

ZHEN-QING CHEN: DEPARTMENT OF MATHEMATICS, UNIVERSITY OF WASHINGTON, SEATTLE, WA 98195, USA, EMAIL: ZQCHEN@UW.EDU

XICHENG ZHANG: SCHOOL OF MATHEMATICS AND STATISTICS, WUHAN UNIVERSITY, WUHAN, HUBEI 430072, P.R.CHINA, EMAIL: XICHENGZHANG@GMAIL.COM