

## Summary for Midterm Two - Math 125

Here is an outline of the material for the second midterm. The core of your studying should be the assigned homework problems: make sure you really understand those well before moving on to other things (like old midterms).

You will definitely be solving a bunch of integrals on this exam (and on the final), so be sure to practice lots of integrals daily. Section 7.5 and the chapter review have lots of extra integrals to practice.

- 6.2 - Volumes (by Cross Section)

- Here is developed the idea that "volume is the integral of (cross sectional) area".
- Although many of the examples we looked at involve *solids of revolution* whose cross-sections are circles, this method applies to any solid that has cross sections whose area can be expressed as a function of  $x$  (or  $y$ ).

- 6.3 - Volume by Cylindrical Shells

- The method of washers/disks is great, but in certain cases we can result in an integral we are unable to evaluate, or, indeed, to setup. So we have another method: the method of cylindrical shells.
- Even if the washer/disk method works, the cylindrical shells method can be easier. Practice will help you decide which method to use on a given solid. Problems 7-14 on page 469 are good for this.

- 6.5 - Average Value of a Function

- This is a very short section. You should understand the definition of the average value of a function on an interval.

- 7.1 - Integration by Parts

- You should understand how to apply the *integration by parts* technique:

$$\int u dv = uv - v \int du$$

- You should be able to recognize integrands for which this technique is particularly appropriate, such as:
  1. a positive integer power of  $x$  times  $\sin x$ ,  $\cos x$ ,  $e^x$  (or related functions)
  2.  $\ln x$  times any power of  $x$
  3.  $e^x$  times  $\sin x$  or  $\cos x$

- Integration by parts can be the method of last resort: if nothing else works, you can almost always try “parts”: if you can differentiate the integrand, you can use it. Whether it helps or not is another question, but it works well with such function as  $\ln x$  and  $\arcsin x$ .

- 7.2 - Trigonometric Integrals

- You should be able to integrate integrals of the form

$$\int \sin^n x \cos^m x dx$$

and

$$\int \tan^n x \sec^m x dx$$

- The strategies for these integrals depend on the *parities* of  $m$  and  $n$  (that is, whether they are even or odd). You should practice all cases. Note that the case where the power of  $\tan x$  is even and the power of  $\sec x$  is odd does not have a clear strategy, so a variety of techniques may be needed.

- 7.3 - Trigonometric Substitution

- This technique exploits trigonometric Pythagorean identities to give useful substitutions that convert quadratic expressions into squares of trig functions. This makes it particularly useful for eliminating square roots (i.e., when you see a quadratic expression inside a square root, there’s a good chance that a trig substitution might be useful).
- One thing that makes this technique more work than a “simple” substitution is the work required to convert back to the original variable once an antiderivative is found. One efficient way to do this is the “triangle method” as described in lecture and the text. Be sure to practice this aspect of this technique.

- 7.4 - Partial Fractions

- This technique is applicable to integrands that are rational functions.
- You should know how to apply this technique to any rational function with a denominator that is factorable as a product of linear factors.
- The idea behind this method is completely algebraic: a rational function whose denominator is a product of linear factors can be expressed as the sum of simpler rational functions, each of which has a denominator which is linear, and a constant numerator. Such simpler functions are easily integrated.
- **Long division of polynomials** is a necessary first step when the numerator has degree equal to or greater than the degree of the denominator. Be sure to practice this if you are not completely comfortable with the method.

- 7.5 - Strategies for Integration

- This section is perhaps the most important. On the exam, the integrals will simply be presented to you for you to solve. You will have to decide which technique, or techniques, to apply to find the antiderivative.

- 7.7 - Numerical Integration

- When we want to evaluate a definite integral, and we cannot find an antiderivative of the integrand, we can approximate the value of the integral. Three techniques we've seen for doing this are

- \* Midpoint Rule
- \* Trapezoid Rule
- \* Simpson's Rule

All three methods are quite similar, and require us to "sample" the function at a number of equally spaced values of  $x$ , then combine these values according to a certain formula which depends on the method.

- 7.8 - Improper Integrals

- There are two types of improper integrals:

1. Those of the form

$$\int_a^{\infty} f(x) dx, \int_{-\infty}^a f(x) dx, \text{ or } \int_{-\infty}^{\infty} f(x) dx$$

2. Those of the form

$$\int_a^b f(x) dx$$

where  $f(x)$  is discontinuous somewhere on the interval  $[a, b]$ .

- We treat improper integrals by defining them as the limits of "proper" integrals. For instance,

$$\int_a^{\infty} f(x) dx = \lim_{k \rightarrow \infty} \int_a^k f(x) dx$$

Be sure to practice your limit taking methods, as these are a key part of evaluating improper integrals.