

M146 Assignment VI Due Thurs May 4, 2006

Overdetermined Systems, Least Squares Fit

Section 43 (page 482) 1, 3, 9, 13, 15

Suppose A is an $m \times n$ matrix and b is a $m \times 1$ column. If $m > n$, the system $Ax = b$ is called overdetermined. There may be no exact solution. The method of best Least Squares Fit is to find x which minimizes the length of $Ax - b$.

- The equations for x are $(A^T \cdot A) \cdot x = A^T \cdot b$, These are called the “Normal Equations.”

(Ex 1) A is a column: $A = \begin{bmatrix} 1 \\ 2 \\ 5 \end{bmatrix}$ and $b = \begin{bmatrix} 6 \\ 7 \\ 8 \end{bmatrix}$. We want the value of x that minimizes the length of the vector $Ax - b$. This will be the value of x that makes the vector $Ax - b$ perpendicular to the vector A . This happens when

$$\begin{aligned} A^T \cdot (Ax - b) &= 0 \\ A^T \cdot A \cdot x - A^T \cdot b &= 0 \\ A^T \cdot A \cdot x &= A^T \cdot b \end{aligned}$$

$$A^T \cdot A = (1 \cdot 1 + 2 \cdot 2 + 5 \cdot 5) = 30; \quad c = A^T \cdot b = (1 \cdot 6 + 2 \cdot 7 + 5 \cdot 8 = 60).$$

The “normal equation” $A^T \cdot A \cdot x = A^T \cdot b$ becomes $30x = 60$; $x = 2$.

Let L be the line through the origin in the direction of $\begin{bmatrix} 1 \\ 2 \\ 5 \end{bmatrix}$. The point on L closest to

$$\begin{bmatrix} 6 \\ 7 \\ 8 \end{bmatrix} \text{ is } \begin{bmatrix} 1 \\ 2 \\ 5 \end{bmatrix} \cdot 2 = \begin{bmatrix} 2 \\ 4 \\ 10 \end{bmatrix}. \text{ The vector of errors is } E = \begin{bmatrix} 1 \\ 2 \\ 5 \end{bmatrix} \cdot 2 - \begin{bmatrix} 6 \\ 7 \\ 8 \end{bmatrix} = \begin{bmatrix} -2 \\ -3 \\ 2 \end{bmatrix}$$

(Ex 2) Find the “best” solution to:

$$\begin{aligned} -x_1 + x_2 &= -2 \\ 0 + 2x_2 &= 6 \\ x_1 + 3x_2 &= 8 \end{aligned}$$

Here A is the 3×2 matrix $A = \begin{bmatrix} -1 & 1 \\ 0 & 2 \\ 1 & 3 \end{bmatrix}$ and $b = \begin{bmatrix} -2 \\ 6 \\ 8 \end{bmatrix}$. We want the vector

$x = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$ that makes the vector $Ax - b$ perpendicular to the plane containing the columns

of A . That is $A^T \cdot (Ax - b) = 0$ which is $A^T \cdot A \cdot x = A^T \cdot b$

$$A^T \cdot A = \begin{bmatrix} -1 & 0 & 1 \\ 1 & 2 & 3 \end{bmatrix} \cdot \begin{bmatrix} -1 & 1 \\ 0 & 2 \\ 1 & 3 \end{bmatrix} = \begin{bmatrix} 2 & 2 \\ 2 & 14 \end{bmatrix}$$

$$\text{and } c = A^T \cdot b = \begin{bmatrix} -1 & 0 & 1 \\ 1 & 2 & 3 \end{bmatrix} \cdot \begin{bmatrix} -2 \\ 6 \\ 8 \end{bmatrix} = \begin{bmatrix} 10 \\ 34 \end{bmatrix}.$$

$$\text{The normal equations are } \begin{bmatrix} 2 & 2 \\ 2 & 14 \end{bmatrix} \cdot \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 10 \\ 34 \end{bmatrix} \quad \text{The solution is } \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 3 \\ 2 \end{bmatrix}$$

$$\text{The vector of errors is } E = \begin{bmatrix} -1 & 1 \\ 0 & 2 \\ 1 & 3 \end{bmatrix} \cdot \begin{bmatrix} 3 \\ 2 \end{bmatrix} - \begin{bmatrix} -2 \\ 6 \\ 8 \end{bmatrix} = \begin{bmatrix} -1 \\ 4 \\ 9 \end{bmatrix} - \begin{bmatrix} -2 \\ 6 \\ 8 \end{bmatrix} = \begin{bmatrix} 1 \\ -2 \\ 1 \end{bmatrix}$$

(Ex 3)) Find the equation of the form $y = b + mt$ which best fits the data:

t	0	2	4
y	2.9	4.2	4.9

We want b and m so that $b + 0m = 2.9$
 $b + 2m = 4.2$. This is an overdetermined system.
 $b + 4m = 4.9$

Here $A = \begin{bmatrix} 1 & 0 \\ 1 & 2 \\ 1 & 4 \end{bmatrix}$, $y = \begin{bmatrix} 2.9 \\ 4.2 \\ 4.9 \end{bmatrix}$. We want $x = \begin{bmatrix} b \\ m \end{bmatrix}$ that minimizes the vector $Ax - y$.

$$A^T \cdot A = \begin{bmatrix} 1 & 1 & 1 \\ 0 & 2 & 4 \end{bmatrix} \cdot \begin{bmatrix} 1 & 0 \\ 1 & 2 \\ 1 & 4 \end{bmatrix} = \begin{bmatrix} 3 & 6 \\ 3 & 20 \end{bmatrix}, \quad \text{and } c = A^T \cdot y = \begin{bmatrix} 1 & 1 & 1 \\ 0 & 2 & 4 \end{bmatrix} \cdot \begin{bmatrix} 2.9 \\ 4.2 \\ 4.9 \end{bmatrix} = \begin{bmatrix} 12 \\ 28 \end{bmatrix}.$$

$$\text{The normal equations are: } \begin{bmatrix} 3 & 6 \\ 3 & 20 \end{bmatrix} \cdot \begin{bmatrix} b \\ m \end{bmatrix} = \begin{bmatrix} 12 \\ 28 \end{bmatrix}. \quad \text{The solution is } \begin{bmatrix} b \\ m \end{bmatrix} = \begin{bmatrix} 3 \\ 0.5 \end{bmatrix}$$

$$\text{The "best" line is } y = 3 + 0.5t. \quad \text{The vector of errors is } E = \begin{bmatrix} 3 \\ 4 \\ 5 \end{bmatrix} - \begin{bmatrix} 2.9 \\ 4.2 \\ 4.9 \end{bmatrix} = \begin{bmatrix} 0.1 \\ -0.2 \\ 0.1 \end{bmatrix}$$

(Ex 4) Find the equation of the form $y = Ae^{kt}$ which best fits the data:

t	1	2	4	7
y	12	14	22	41

Take logarithms of the y values to get

t	1	2	4	7
w	2.48	2.64	3.09	3.70

We want the line $w = b + mt$ which best fits this data.

We want b and m so that

$$\begin{aligned} b + 1m &= 2.48 \\ b + 2m &= 2.64 \\ b + 4m &= 3.09 \\ b + 7m &= 3.70 \end{aligned}, \quad \text{which is overdetermined.}$$

Here $A = \begin{bmatrix} 1 & 1 \\ 1 & 2 \\ 1 & 4 \\ 1 & 7 \end{bmatrix}$, $w = \begin{bmatrix} 2.48 \\ 2.64 \\ 3.09 \\ 3.70 \end{bmatrix}$. Want $x = \begin{bmatrix} b \\ m \end{bmatrix}$ that minimizes the vector $Ax - w$.

$$A^T \cdot A = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 0 & 2 & 4 & 7 \end{bmatrix} \cdot \begin{bmatrix} 1 & 1 \\ 1 & 2 \\ 1 & 4 \\ 1 & 7 \end{bmatrix} = \begin{bmatrix} 4 & 14 \\ 14 & 70 \end{bmatrix}, \quad \text{and}$$

$$c = A^T \cdot w = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & 2 & 4 & 7 \end{bmatrix} \cdot \begin{bmatrix} 2.48 \\ 2.64 \\ 3.09 \\ 3.70 \end{bmatrix} = \begin{bmatrix} 11.91 \\ 46.02 \end{bmatrix}.$$

The normal equations are $\begin{bmatrix} 4 & 14 \\ 14 & 70 \end{bmatrix} \cdot \begin{bmatrix} b \\ m \end{bmatrix} = \begin{bmatrix} 11.9 \\ 46.0 \end{bmatrix}$

The solution is $\begin{bmatrix} b \\ m \end{bmatrix} = \begin{bmatrix} 2.3 \\ 0.2 \end{bmatrix}$, (numbers rounded off).

The “best” line is $w = 2.3 + 0.2t$ The exponential is $y = e^{2.3}e^{0.2t} = 10e^{0.2t}$

M146 Sample Quiz #06 for Thursday, May 4, 2006

(1) Let $C = \begin{bmatrix} 1 \\ 2 \\ 5 \end{bmatrix}$. Find the point on the line $x = Ct$ closest to the point $b = \begin{bmatrix} 3 \\ 4 \\ 7 \end{bmatrix}$.

(2) $A = \begin{bmatrix} 1 & 3 \\ 2 & 2 \\ 5 & 1 \end{bmatrix}$; $b = \begin{bmatrix} 3 \\ 4 \\ 7 \end{bmatrix}$. Find the best (least squares) solution to $Ax = b$

(3) The following data were obtained for the length y in centimeters of a human fetus versus the age t in weeks. Find a linear function $y = b + mt$ which best fits the data.

Age t	12	20	28	40
Length y	10	25	38	53

(4) Radioactive sample Y is decaying exponentially. The data shows the amount (y in grams) of Y at times (t in days). Find an exponential function $y = Ae^{rt}$ which best fits the data.

t	0	1	2	3	4	5
y	100	82	67	55	45	37

(5) The table shows the length-weight relation for Pacific halibut. Find an power function $W = \alpha L^\beta$ which best fits the data.

L	0.5	1.0	1.5	2.0	2.5
W	1.3	10.4	35	82	163

(6) The velocity of an enzymatic reaction with Michaelis-Menton kinetics is given by

$$v(s) = \frac{\alpha s}{1 + \beta s}$$

Inverting this gives the Lineweaver-Burke equation:

$$\frac{1}{v} = \frac{1}{\alpha} \frac{1}{s} + \frac{\beta}{\alpha}$$

With the change of variables $\frac{1}{v} = w$ and $\frac{1}{s} = u$, this becomes:

$$w = \frac{1}{\alpha} u + \frac{\beta}{\alpha}$$

Find the Michaelis-Menton equation which best fits the data:

s	1	2.5	5	10	20
v	4.1	6.1	9.3	12.9	17.1