

Math 125, Sections C and F, Fall 2014, Midterm I

October 16, 2014

Name Solutions

TA/Section _____

Instructions.

- There are 4 questions. The exam is out of 40 points.
- You are allowed to use one page of notes written only on one side of the sheet in your own handwriting. Hand in you note sheet with your exam.
- Calculators are NOT allowed. Put away ALL electronic devices.
- For your integrals you may use the following formulas. Anything else must be justified by your work.

$$\int x^n dx = \frac{x^{n+1}}{n+1} + C, n \neq -1 \quad \int e^x dx = e^x + C \quad \int \frac{1}{x} dx = \ln|x| + C$$

$$\int \sin x dx = -\cos x + C \quad \int \cos x dx = \sin x + C \quad \int \sec^2 x dx = \tan x + C$$

$$\int \csc x \cot x dx = -\csc x + C \quad \int \sec x \tan x dx = \sec x + C = \int \csc^2 x dx = -\cot x + C$$

$$\int \frac{1}{\sqrt{1-x^2}} dx = \sin^{-1} x + C \quad \int \frac{1}{1+x^2} dx = \tan^{-1} x + C$$

- **Show your work.** If I cannot read or follow your work, I cannot grade it. You may not get full credit for a right answer if your answer is not justified by your work. If you continue at the back of a page, make a note for me.

Question	points
1	
2	
3	
4	
Total	

Solve

1. (10 points) Evaluate the following integrals.

(a) $\int 7 \cos(\theta) \sin^2(\theta) d\theta$

$$u = \sin \theta$$
$$du = \cos \theta d\theta$$

$$\int 7 u^2 d\theta = \frac{7}{3} u^3 + C = \frac{7}{3} \sin^3 \theta + C$$

(b) $\int_0^1 \frac{x}{1+5x} dx$


$$u = 1 + 5x \rightarrow x = \frac{u-1}{5}$$
$$du = 5 dx$$

$$= \int_1^6 \frac{\frac{u-1}{5}}{u} \cdot \frac{1}{5} du = \frac{1}{25} \int_1^6 \frac{u-1}{u} du = \frac{1}{25} \int_1^6 \left(1 - \frac{1}{u}\right) du$$

$$= \frac{1}{25} [u - \ln|u|]_1^6 = \frac{1}{25} (5 - \ln 6)$$

(c) $\int_{-1}^1 x e^{x^8} dx = 0$ because $x e^{x^8}$ is an odd function.

2. (10 points) Define $g(x) = \int_4^x f(t)dt$ where the graph of $f(t)$ is given below.



(a) Evaluate the following:

$$g(0) = \int_0^0 f(t)dt = - \int_0^4 f(t)dt = -40$$

$$g(4) = 40$$

$$g(16) = \frac{2 \cdot 10}{2} - \frac{6 \cdot 10}{2} + \frac{4 \cdot 10}{2} = 10 - 30 + 20 = 0$$

$$g'(17) = f(17) = 10$$

$$g''(11) = f'(11) = \frac{20}{8} = 2.5$$

(b) Express $g(22) - g(18)$ as a definite integral and estimate it with $n = 4$ and leftpoints. This question will be graded with a reasonable allowance for estimation error.

$$\int_{18}^{22} f(t)dt \approx [f(18) + f(19) + f(20) + f(21)] \Delta t$$

$$\approx 10 + 9.8 + 8.8 + 5.8$$

$$\approx 34.4$$

$$\Delta t = \frac{22-18}{4} = 1$$

(c) If $h(x) = \int_4^{x^3} f(t)dt$, what is $h'(2)$?


$$h(x) = g(x^3)$$

$$h'(x) = g'(x^3) \cdot 3x^2$$

$$= f(x^3) \cdot 3x^2$$

$$h'(2) = f(8) \cdot 3 \cdot 4 = 0 \quad \text{120}$$

3. Find the area of the region shown below bounded by the curve $x = -y^3 + y^2$ and the line $x = -2y$.



Points of intersection:

$$-y^3 + y^2 = -2y$$

$$-y^3 + y^2 + 2y = 0$$

$$-y(y^2 - y - 2) = 0$$

$$-y(y-2)(y+1) = 0$$

$$y=0 \quad y=2 \quad y=-1$$

$$x=0 \quad x=-4 \quad x=2$$

$$A = \int_0^2 (-y^3 + y^2) - (-2y) dy$$


$$= \int_0^2 -y^3 + y^2 + 2y dy = \left[-\frac{y^4}{4} + \frac{y^3}{3} + y^2 \right]_0^2$$

$$= -4 + \frac{8}{3} + 4 = \frac{8}{3}$$

4. (11 points)

- (a) Sketch the region between the parabola $y = x^2 - 12x + 32$ and the line $y = 3x - 4$. Label all intersection points.

$$y = x^2 - 12x + 32 = (x-8)(x-4)$$



$$3x - 4 = x^2 - 12x + 32$$

$$0 = x^2 - 15x + 36$$

$$0 = (x-12)(x-3)$$

$$\begin{array}{l} x=12 \\ y=32 \end{array} \quad \begin{array}{l} x=3 \\ y=5 \end{array}$$

$$x-1$$

$$y = (x-9)(x-5) = x^2 - 14x + 45$$

$$y = 3(x-1) - 4 = 3x - 7$$

- (b) Set up an integral to calculate the volume of the solid formed by rotating this region about the y -axis. Do NOT integrate.

$$\int_3^{12} 2\pi x [3x - 4 - (x^2 - 12x + 32)] dx$$

- (c) Set up an integral to calculate the volume of the solid formed by rotating this region about the horizontal line $y = 40$. Do NOT integrate.

$$\int_3^{12} \pi \left[(40 - (x^2 - 12x + 32))^2 - (40 - (3x - 4))^2 \right] dx$$

- (d) Set up an integral to calculate the volume of the solid formed by rotating this region about the vertical line $x = 4$. Do NOT integrate.

This is rotating the part to the left of $x=4$ about $x=4$

$$\int_4^{12} 2\pi (x-4) (3x-4 - (x^2 - 12x + 32)) dx$$