Name:

Student Number:

No books or notes allowed on this exam.

Find the absolute maximum and absolute minimum of

$$f(x) = x - 2 \arctan x$$

on the interval [0,4]. Use sentences to justify your answer (don't just circle a number, but use the reasoning we learned in class.)

Solution: [J. Stewart, Page 278] <u>The Closed Interval Method</u> To find the *absolute* maximum and minimum values of a continuous function f on a closed interval [a, b]:

1. Find the values of f at the critical numbers of f in (a,b). Set f'(x) = 0.

Solve for the critical numbers:

$$f'(x) = 1 - 2 \cdot \left(\frac{1}{1+x^2}\right) = 0$$

$$1 = \frac{2}{1+x^2}$$

$$1 + x^2 = 2$$

$$x^2 = 1 \Longrightarrow x = \pm 1.$$

Since -1 is not in the domain, the interval [0,4], we have x=1 as our critical number. Plug in x=1 in f. We have

$$f(1) = 1 - 2 \arctan 1 = 1 - 2 \cdot \frac{\pi}{4} = 1 - \frac{\pi}{2}.$$

2. Find the values of f at the endpoints of the interval.

$$f(0) = 0 - 2 \arctan 0 = 0.$$

 $f(4) = 4 - 2 \arctan 4.$ (No need to simplify further.)

3. The largest of the values from Steps 1 and 2 is the absolute maximum value; the smallest of these values is the absolute minimum value.

To get the credit in this part, you only need to state the step in sentences. However, there is a way to find the largest value and the smallest value without using the calculator. Notice that $1 - \frac{\pi}{2}$ is negative because π is bigger than 2. Also, $4 - 2 \arctan 4$ is positive because $2 \arctan 4$ is less than π . Thus,

$$1 - \frac{\pi}{2} < 0 < 4 - 2\arctan 4.$$

Therefore, f(1) is the absolute minimum and f(4) is the absolute maximum.