


F.I.L.C.T.

the


FUNNY LITTLE CALCULUS TEXT



VOLUME 1:
SINGLE
VARIABLE


by
ROBERT CHRIST

the Andrea Mitchell
University Professor of
Mathematics & Electrical /
Systems Engineering at
the University of Pennsylvania



What is this book?

It's a Calculus text. It's little. It's funny.
It's not a replacement for your
standard, problemfull text.



This text is meant to be read and enjoyed.

PREREQUISITES



This text assumes you've seen some Calculus before: you know what to do [differentiate/integrate] and how to do it, but you don't know what it really means -- like everything else in Life...  

INSTRUCTIONS

This is for the "big picture" (~~some little pictures too...~~). You should find a good source for homework problems, details, & proofs.

BASICALLY

- 10 READ
- 20 THINK
- 30 SOLVE PROBLEMS
- 40 GOTO 10

AND THEY ALL SAD... 

THE COLORS! IT'S TOO MUCH!! 
not to mention the pretentious literary allusions commingled with crude humor 
Are you, in fact, Barking Mad? 

THANKS!


To my students, for teaching me...

This book was written on a Fujitsu tablet PC using Microsoft Journal.




TABLE of CONTENTS


CHAPTER 1: FUNCTIONS




CHAPTER 2: DIFFERENTIATION




CHAPTER 3: INTEGRATION




CHAPTER 4: APPLICATIONS




CHAPTER 5: DISCRETIZATION



This is single-variable Calculus: stay tuned for Volume II




MULTIPASS!



~~MULTIVARIABLE~~

FLCT Chapter 1:

functions



You can't get far in Calculus without understanding functions:
They are the DNA of Mathematics... 
that is, elementary building blocks...

FUNCTIONS

This is a function.

x = INPUT

$f(x)$ = OUTPUT

DOMAIN

x



function machine

f

RANGE
here, here or the

$f(x)$

These may or may not be numbers... but to be a function:

SAME INPUT \rightarrow SAME OUTPUT

EXAMPLES

$f(x) = x^2$ is a function

$f(x) = \tan^{-1}(x)$ is not a function many angles have $\tan^{-1} x$, so we use "ARCTAN" to fix range...

$f(\text{location, time}) = (\text{temperature, windspeed})$ is a function

1 INPUT

1 OUTPUT

MULTIVARIATE

$f(\text{hardwork}) = \text{success}$ is not a function (success depends on more than one variable...)

OPERATIONS

You can do the typical things to typical functions:




MY DEAR AUNT SALLY

The most important operations are the following:

COMPOSITION


$$(f \circ g)(x) = f(g(x))$$




INVERSE

$$(f^{-1} \circ f)(x) = x = (f \circ f^{-1})(x)$$

FOR ALL x



EXAMPLE $\sqrt[3]{x}$ is the inverse of x^3
 \sqrt{x} [positive square root] is the inverse of x^2 $(x \geq 0)$
 a^x and $\ln_a x$ are inverses



POLYNOMIALS & POWERS

A **MONOMIAL** is a function of the form: ax^n (let's say $n = 0, 1, 2, \dots$)

A **POLYNOMIAL** is $p(x) = \sum_{k=0}^n a_k x^k = a_0 + a_1 x + a_2 x^2 + \dots + a_n x^n$ **TERMS** **DEGREE**

A **RATIONAL** function is a quotient $r(x) = \frac{p(x)}{q(x)}$ **Polys**

Let's think... what is... **CALCULATING EXISTENTIAL ANGST...**

x^0 : EASY. $x^0 = 1$ for all x

 x^{-1} : EASY. $x^{-1} = \frac{1}{x}$ for all $x \neq 0$

x^{-n} : EASY. $x^{-n} \cdot x^n = x^{-n+n} = x^0 = 1 \rightarrow x^{-n} = \frac{1}{x^n}$

$x^{\frac{1}{n}}$: EASY. $(x^{\frac{1}{n}})^n = x^{\frac{n}{n}} = x \rightarrow x^{\frac{1}{n}} = \sqrt[n]{x}$

But: $x^{\sqrt{2}} = ?$ $x^{\pi} = ?$ $x^{\sqrt[3]{3}} = ?$ What are these?

If you try to compute 2^π , you might approximate...

$$2^\pi \approx 2^{\frac{314}{100}} = \sqrt[100]{2^{314}}$$



A certain limiting process seems inevitable. The principled approach to such phenomena is encapsulated in...

THE EXPONENTIAL FUNCTION

The most important function in Calculus is

$$\text{EXP}(x) := e^x$$

where $e = 2.718281828459045235360287471352662497779787848$

and x is, well, x
actually, e is just



But there is a better, subtler definition:

$$\text{SERIES } e^x = 1 + x + \frac{1}{2}x^2 + \frac{1}{6}x^3 + \frac{1}{24}x^4 + \frac{1}{120}x^5 + \frac{1}{720}x^6 + \dots$$


AD INFINITUM

DO NOT $\sum_{k=0}^{\infty} \frac{1}{k!} x^k$ **FEAR Sigma - "SUM"**

Where $k! = k(k-1)(k-2) \dots (3)(2)(1)$ and $0! = 1$ by convention

! \Rightarrow "FACTORIAL" oh yeah?
well, I'm 2.


This is the definition of e^x : it requires an INFINITE SERIES




Although this looks complicated, it only takes a few terms to get a decent approximation. For the time being, pretend that such infinite sums are polynomials. I promise no harm will come to you in so doing (for now...).

SPECIAL FUNCTIONS


LOGS



= "natural log" = inverse of e^x



TRIGS



... & inverse of the other trigs, like ARCCOS, ARCTAN, etc., and the other trigs: e.g., ARCSIN, COS, TAN, SEC, CSC, ...

HYPERSOULC TRIGS




I must put it in my head!

$$\sinh x = \frac{e^x - e^{-x}}{2}$$

$$\cosh x = \frac{e^x + e^{-x}}{2}$$

$$\tanh x = \frac{\sinh x}{\cosh x}$$



SECH CSC COTH
NOT

EULER'S FORMULA


NOT "yew-lors" but "oi-lers": it's a Swiss thing!

The special functions you know & love are all connected

$$\cos^2 x + \sin^2 x = 1$$


$$\cosh^2 x - \sinh^2 x = 1$$

$$\cos x + i \sin x = e^{ix}$$




RECALL:
 $i = \sqrt{-1}$
 $i^2 = -1$

These & similar formulae are illuminative



The ability to switch between algebraic and geometric thinking is crucial...



SPECIAL SERIES

We know...

$$e^x = 1 + x + \frac{1}{2!}x^2 + \frac{1}{3!}x^3 + \frac{1}{4!}x^4 + \frac{1}{5!}x^5 + \dots$$

Let us see what happens to other special functions:

$$\sinh x = \frac{1}{2}e^x - \frac{1}{2}e^{-x} \quad \text{EXPAND } e^x \text{ & } e^{-x}!$$



$$\begin{aligned} &= \frac{1}{2}(1 + x + \frac{1}{2!}x^2 + \frac{1}{3!}x^3 + \frac{1}{4!}x^4 + \frac{1}{5!}x^5 + \dots) \\ &\quad - \frac{1}{2}(1 + (-x) + \frac{1}{2!}(-x)^2 + \frac{1}{3!}(-x)^3 + \frac{1}{4!}(-x)^4 + \frac{1}{5!}(-x)^5 + \dots) \end{aligned}$$

$$= \cancel{1} + x + \cancel{0}x^2 + \frac{1}{3!}x^3 + \cancel{0}x^4 + \frac{1}{5!}x^5 + \dots$$

$$\begin{aligned} \text{likewise... } \sinh x &= x + \frac{1}{3!}x^3 + \frac{1}{5!}x^5 + \dots &= \sum_{k=0}^{\infty} \frac{1}{(2k+1)!} x^{2k+1} & \text{How odd!} \\ \cosh x &= 1 + \frac{1}{2!}x^2 + \frac{1}{4!}x^4 + \dots &= \sum_{k=0}^{\infty} \frac{1}{(2k)!} x^{2k} & \text{How even!} \end{aligned}$$

Euler's formula gives us a series formula for \sin/\cos

$$\begin{aligned}
 e^{ix} &= 1 + ix + \frac{1}{2!}(ix)^2 + \frac{1}{3!}(ix)^3 + \frac{1}{4!}(ix)^4 + \frac{1}{5!}(ix)^5 + \dots \\
 &= 1 + ix - \frac{1}{2!}x^2 - i\frac{1}{3!}x^3 + \frac{1}{4!}x^4 + i\frac{1}{5!}x^5 - \dots \\
 &= \left(1 - \frac{1}{2!}x^2 + \frac{1}{4!}x^4 - \frac{1}{6!}x^6 + \dots\right) + i\left(x - \frac{1}{3!}x^3 + \frac{1}{5!}x^5 - \frac{1}{7!}x^7 + \dots\right) \\
 &\stackrel{\text{EULER}}{=} \cos(x) + i \sin(x)
 \end{aligned}$$

COMPLEX! REAL IMAGINARY

$\cos(x) = 1 - \frac{1}{2!}x^2 + \frac{1}{4!}x^4 - \dots = \sum_{k=0}^{\infty} \frac{(-1)^k}{(2k)!} x^{2k}$ TRICKY...

$\sin(x) = x - \frac{1}{3!}x^3 + \frac{1}{5!}x^5 - \dots = \sum_{k=0}^{\infty} \frac{(-1)^k}{(2k+1)!} x^{2k+1}$



Other special series come with RULES

$$1/1/1/1.1'1/1/1/1/1.$$

$$1 + \frac{1}{2} + \frac{1}{4} + \frac{1}{8} + \frac{1}{16} + \frac{1}{32} + \dots = 2$$

$$\sum_{k=0}^{\infty} \left(\frac{1}{2}\right)^k = 2$$

This is a special case of the GEOMETRIC SERIES:

$$\frac{1}{1-x} = 1 + x + x^2 + x^3 + x^4 + \dots = \sum_{k=0}^{\infty} x^k$$

(proof: $(1-x)(1+x+x^2+x^3+\dots) = (1+x+x^2+x^3+\dots) - (x+x^2+x^3+x^4+\dots) = 1$ voila!) but does $1+2+4+8+16+\dots = \frac{1}{1-x} = -1$? No! Remember $|x|<1$!

BUT! ONLY
FOR $|x|<1$
unless you are a physicist

TAYLOR SERIES


It seems as though many standard functions may have infinite series expressions. This leads to the first big idea of FLCT:

ANY "REASONABLE" $f(x)$ EQUALS —

$\sum_{k=0}^{\infty} c_k x^k$ FOR SOME c_k CONSTANTS —

Such a series expansion is called a TAYLOR SERIES about $x=0$. It means that, near 0, $f(x)$ looks like a polynomial... (this also goes by the name "MACLAURIN" SERIES)

It's best to think in terms of POLYNOMIAL APPROXIMATION. You know that the tangent line to $y = e^x$ at $(0, 1)$ is $1+x$. The best fit quadratic function is $1+x+\frac{1}{2}x^2$. Keep going: the TAYLOR POLYNOMIAL to e^x at $x=0$ of degree n is:

$$1 + x + \frac{1}{2!}x^2 + \dots + \frac{1}{n!}x^n$$


To get a tangent line to any $f(x)$ at $x=0$, you use $f'(0)$, the derivative. It should not surprise you to learn that the other polynomial approximations to $f(x)$ are also based on derivatives. The TAYLOR SERIES of $f(x)$ about $x=0$ is:

$$\sum_{k=0}^{\infty} \frac{f^{(k)}(0)}{k!} x^k = f(0) + f'(0)x + \frac{1}{2!} f''(0)x^2 + \frac{1}{3!} f'''(0)x^3 + \dots$$

+ going up


means " k "th derivative"

$$= f(x) \quad \text{for most functions, at least}$$



By changing coordinates $x \mapsto x-a$ we obtain the TAYLOR SERIES of $f(x)$ ABOUT $x=a$:

$$\sum_{k=0}^{\infty} \frac{f^{(k)}(a)}{k!} (x-a)^k = f(a) + f'(a)(x-a) + \frac{1}{2!} f''(a)(x-a)^2 + \frac{1}{3!} f'''(a)(x-a)^3 + \dots$$



For x near a , a truncation of this to a polynomial gives the best polynomial approximation to f (near a).

COMPUTING TAYLOR SERIES

Since we all know how to differentiate like a superhero, it's no big deal to compute a few examples:

EXAMPLE $\frac{d}{dx}(e^x) = e^x \Rightarrow (e^x)^{(k)}(0) = e^0 = 1$ Thus,

$$e^x = 1 + \frac{1}{1!}x + \frac{1}{2!}x^2 + \frac{1}{3!}x^3 + \dots \quad \text{AHA!}$$

BUT... ABOUT $x=10$, THIS IS A BAD APPROXIMATION: TRY...


$$e^x = e^{10} + \frac{e^{10}}{1!}(x-10) + \frac{e^{10}}{2!}(x-10)^2 + \frac{e^{10}}{3!}(x-10)^3 + \dots = e^{10} \cdot e^{x-10}$$



OF COURSE

EXAMPLE 5

$$\begin{array}{ccccccc} \sin x & \xrightarrow{\frac{d}{dx}} & \cos x & \xrightarrow{\frac{d}{dx}} & -\sin x & \xrightarrow{\frac{d}{dx}} & -\cos x \xrightarrow{\frac{d}{dx}} \sin x \\ \downarrow x=0 & & \downarrow x=0 & & \downarrow x=0 & & \downarrow x=0 \\ 0 & & 1 & & 0 & & -1 \end{array}$$



So... $\sin x = 0 + \frac{1}{1!}x + \frac{0}{2!}x^2 - \frac{1}{3!}x^3 + \frac{0}{4!}x^4 + \dots = x - \frac{x^3}{3!} + \frac{x^5}{5!} - \frac{x^7}{7!} + \dots$

But, in general, computing lots of derivatives is NO FUN. Fortunately, pretending that Taylor series are polynomials allows one to perform computations like these:

EXAMPLE 6 $e^{\sin(x^2)} = e^{(x^2 - \frac{x^6}{3!} + \frac{x^{10}}{5!} - \frac{x^{14}}{7!} + \dots)} = 1 + (x^2 - \frac{x^6}{3!} + \frac{x^{10}}{5!} - \dots) + \frac{1}{2}(x^2 - \frac{x^6}{3!} + \dots)^2 + \frac{1}{3!}(x^2 + \dots)$

$= 1 + x^2 + \frac{1}{2}x^4 + \frac{1}{24}x^8 + \text{H.O.T.}$ i.e., stuff I don't care about...

This requires ALGEBRA , but only polynomial algebra.

EXAMPLE 7 $\frac{1}{1-x} = 1 + x + x^2 + x^3 + x^4 + \dots + x^n + \dots$ GEOMETRIC SERIES $|x| < 1$!

SUBS

IS

THIS

LEGIT

$$\begin{aligned} \frac{1}{1+x} &= 1 - x + x^2 - x^3 + x^4 + \dots + (-1)^n x^n + \dots = \sum_{n=0}^{\infty} (-1)^n x^n \\ \frac{1}{(1-x)^2} &= 1 + 2x + 3x^2 + 4x^3 + \dots + nx^n + \dots = \sum_{n=0}^{\infty} nx^n \\ \ln(1+x) &= x - \frac{x^2}{2} + \frac{x^3}{3} - \frac{x^4}{4} + \dots + (-1)^{n-1} \frac{x^n}{n} = \sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{n} x^n \\ \ln x &= (x-1) - \frac{(x-1)^2}{2} + \frac{(x-1)^3}{3} + \dots + (-1)^{n-1} \frac{(x-1)^n}{n} = \sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{n} (x-1)^n \end{aligned}$$

MIRABILE DICTU, you can pull this off. No 'blemo, as long as you remember (in this case) $|x| < 1$.

(= Latin: "cool dat. word!"



"CALCULVM VIRVMQVE CANO"

AND NOW THE BEGINNING


The time has come to do Calculus. We begin at the beginning with the notion of LIMITS. Perhaps you recall ...

$\lim_{x \rightarrow a} f(x) = L \iff$ For every $\epsilon > 0$, there exists a $\delta > 0$ so that whenever $0 < |x - a| < \delta$ one has $|f(x) - L| < \epsilon$.

You have an intuition for what this means, but it is worth learning to parse the definition.

HINTS: δ = TOLERANCE on INPUT

ϵ = TOLERANCE on OUTPUT



This is like an adversarial game.

Player 1 picks ϵ ; Player 2 must choose a sufficiently small $\delta > 0$. Engineers/scientists consistently use tolerances for error: same deal.

It is a slight generalization to define $\lim_{x \rightarrow \infty} f(x)$ and $\lim_{x \rightarrow -\infty} f(x)$. Note that here, as ever, existence is not automatic. Our motivation for investigating limits is to define

continuity

A continuous function is one for which (INSERT PENCIL METAPHOR)

DEF $\lim_{x \rightarrow a} f(x) = f(a) \quad \forall a$ \forall = "for all" (not to be confused with \otimes = "for Me")

In particular, the limits (2-SIDED) must exist! Most functions you see in real life are continuous: polynomials, exponentials, sines, cosines, etc.; beware SEC, CSC, LN and rational functions. Certain digital signals are discontinuous. But no biggie...

The end of this Chapter has arrived, in which the hard work of understanding functions as Taylor series pays off...




LIMITS & SERIES

EXAMPLE

Who remembers this one?

$$\lim_{x \rightarrow 0} \frac{\sin x}{x} = 1$$



THE REASON:

$$\lim_{x \rightarrow 0} \frac{\sin x}{x} = \lim_{x \rightarrow 0} \frac{x - \frac{x^3}{3!} + \text{HOT}}{x} = \lim_{x \rightarrow 0} 1 - \frac{x^2}{3!} + \text{HOT} = 1$$

LOTS OF x
CANCEL!



THE TAYLOR SERIES ABOUT $x=0$ IS $\cdot 1 + \text{HOT} \dots$

"AHA!", you say, "I could have used l'Hôpital's rule." Well guess what: I just used something better: l'Hôpital's WHOLEGOSH DARNED Raison d'être!

HULK SMASH!

If all you know is THIS RULE and THAT RULE and TRICKS IN MY HEAD, then you don't know Calculus -- you are just a PATTERNMATCHINGAUTOMAZOMBIE!



Don't. Be. That. Way. Learn the reasons why the rules/tools work!

Why is it that l'Hôpital's rule doesn't always work? Or, if it doesn't, why you keep on taking derivatives?


Assume: $\lim_{x \rightarrow a} f(x) = 0 = \lim_{x \rightarrow a} g(x)$

THEN:

$$\text{Expand about } x=a \quad \lim_{x \rightarrow a} \frac{f(x)}{g(x)} = \lim_{x \rightarrow a} \frac{f(a) + f'(a)(x-a) + \frac{1}{2}f''(a)(x-a)^2 + \text{HOT}}{g(a) + g'(a)(x-a) + \frac{1}{2}g''(a)(x-a)^2 + \text{HOT}}$$

Lots of $(x-a)$'s TO FACTOR-

This is why l'Hôpital's rule is.



EXAMPLE

$$\lim_{x \rightarrow 1} \frac{1 - \cos(x-1)}{\ln^2 x} = \lim_{x \rightarrow 1} \frac{1 - \left(1 - \frac{(x-1)^2}{2!} + \text{HOT}\right)}{\left((x-1) - \frac{(x-1)^2}{2} + \text{HOT}\right)^2} = \lim_{x \rightarrow 1} \frac{\frac{1}{2!}(x-1)^2 + \text{HOT}}{(x-1)^2 + \text{HOT}}$$

(x-1) FACTORS
CANCEL!

You need the 2nd order term to compute the limit.

ORDERS OF GROWTH

Calculus requires a familiarity with functional growth. We have used H.O.T. for terms such as $(x-a)^{\text{big}}$, which go to zero quickly as $x \rightarrow a$. How quickly? What if $x \rightarrow \infty$?

CONSIDER:

$$\lim_{x \rightarrow \infty} \frac{2x^2 - 5x + 3}{3x^2 - 11} = \lim_{x \rightarrow \infty} \frac{2 - \frac{5}{x} + \frac{3}{x^2}}{3 - \frac{11}{x^2}} = \frac{2}{3}$$

DOMINANT TERMS COEFFICIENTS

$$\text{or... } \lim_{x \rightarrow \infty} \frac{e^x}{x^2} = \lim_{x \rightarrow \infty} \frac{1 + x + \frac{1}{2!}x^2 + \frac{1}{3!}x^3 + \dots}{x^2} = \lim_{x \rightarrow \infty} \frac{\frac{1}{x^2} + \frac{1}{x} + \frac{1}{2!} + \frac{x}{3!} + \dots}{1} = +\infty$$



This line of reasoning shows that $\lim_{x \rightarrow \infty} \frac{e^x}{x^n} = \infty$ for all $n \geq 0$

EXPONENTIALS BEAT POLYNOMIALS

There is a hierarchy of functions based on ASYMPTOTICS: It pays to know who is on their way to the top at what speed.




FACTORIAL GROWTH



EXPONENTIAL GROWTH



POLYNOMIAL GROWTH



LOGARITHMIC GROWTH



NO GROWTH

x^x or $x!$

(Protein conformations;
Ways to live your life -- assuming no fate)

e^x or $\cosh x$ or $\sinh x$

(Compound investment
or credit card debt
or Moore's Law)

x^n $n > 0$

(area/volume growth;
network utility;
falling bodies)

$\ln x$

(crypto key length;
Richter scale; pH)

CONST or SIN

(mess on my desk)

In the next chapter, we continue with derivatives...



of course it's
small & local

+ + + + +
+ + + + +
WHOA...
+ + + + +


Don't try this with substances
is fatal as x
+ + + + +

the subject of asymptotic growth keeps going: look up big O notation on Wikipedia

FLCT.


~~Chapter 2:~~

~~Differentiation~~






So you aced Calculus in high school and can differentiate while unconscious. Well that's great. But do you know what it all means?



SLOPE?
NOPE.

DEFINITION

There is a distinction between definition & interpretation: this is what a derivative is



$$\left. \frac{df}{dx} \right|_{x=a} = f'(a) = \lim_{h \rightarrow 0} \frac{f(a+h) - f(a)}{h}$$

ONLY FOR SINGLE-VARIABLE FUNCTIONS!



Like most definitions, this one is quickly assimilated & forgotten... I assume you've been there, done that.

The computation of derivatives is usually easy, based on a few simple rules: first, and most important is the overwhelming question


=WHAT DOES A DERIVATIVE MEAN?

(As with other things in life, interpretation matters as much as definition.)

NOTATION ALERT!		
f'	$\frac{df}{dx}$	OK!
$\frac{df}{dx}$	$\frac{df}{dx}$	NOT OK
$\frac{df}{dx}$	$\frac{df}{dx}$	DON'T GO THERE

APPROXIMATION

In keeping with the theme of **FLCT**, we begin with a Taylor series interpretation. This is, admittedly circular logic, but, in the words of the great Archimedes, "My *YOU TOUR MY LOGIC TOWER!*"



what does $f(a+h)$ look like for h small?

=TAYLOR EXPAND $f(x)$ ABOUT $x=a$ =

$$f(x) = f(a) + f'(a)(x-a) + \frac{1}{2}f''(a)(x-a)^2 + \text{HOT.}$$

LINEAR APPROXIMATION ERROR TERMS



Consider the substitution: $x = a + h$


$$f(a+h) = f(a) + \underbrace{f'(a)h}_{\text{LEADING TERM}} + \underbrace{\frac{1}{2}f''(a)h^2 + \frac{1}{6}f'''(a)h^3}_{\text{VERY SMALL FOR } h \approx 0} + \text{HOT}$$

from which follows a tautological computation:

MATH GURU!

$$\begin{aligned} f'(a) &= \lim_{h \rightarrow 0} \frac{1}{h} (f(a+h) - f(a)) \quad (\text{substitute/simplify}) \\ &= \lim_{h \rightarrow 0} \frac{1}{h} (f'(a)h + \frac{1}{2}f''(a)h^2 + \text{HOT}) = \lim_{h \rightarrow 0} (f'(a) + \frac{1}{2}f''(a)h + \text{HOT}) \\ &= f'(a) \end{aligned}$$

SO WHAT? If you think about it...



The derivative of f at a is the coefficient of the 1^{st} ORDER TERM in the polynomial approximation to $f(a+h)$


If you didn't know the definition of the derivative, you could construct it from the Taylor series...

EXAMPLE What is the area of a square of side length $4\frac{1}{7}$?

This is easily approximated without a calculator...

Let $x = \text{SIDE LENGTH}$; $f(x) = x^2$ AREA

Expanding about $a = 4$ (a nice number)...




$$f(x) = f(4) + f'(4)(x-4) + \text{HOT}$$

$$= 16 + 8'(\frac{1}{7}) + \text{HOT}$$

$$\approx 17\frac{1}{7}$$


To FIRST ORDER



The higher order term (singular) is $\frac{1}{4}h^2 = (\frac{1}{7})^2$

you can "see" the terms of a Taylor series in such examples, with the derivative as the first-order term

RATE OF CHANGE
function-mechanical...



Perhaps the best interpretation of $f'(a)$ is

$$f'(a) = \lim_{x \rightarrow a} \frac{f(x) - f(a)}{x - a} \quad \begin{matrix} \leftarrow \Delta \text{ OUTPUT} \\ \leftarrow \Delta \text{ INPUT} \end{matrix}$$

(PROOF: let $x = a+h$; then $\Delta = \lim_{h \rightarrow 0} \frac{f(a+h) - f(a)}{(a+h) - a}$)

thus, the derivative is a RATE of change



The derivative does not tell you the actual change in the output...
JUST THE RATE OF CHANGE!

~~rate~~ **RATES**

VELOCITY: if $x(t)$ = position of an object at time t , then the derivative $v(t) = \frac{dx}{dt}$ is the VELOCITY, where $v > 0$ means \rightarrow and $v < 0$ means \leftarrow (RECALL: VELOCITY = SPEED + DIRECTION)


ACCELERATION: in the above, $a = \frac{dv}{dt} = \frac{d^2x}{dt^2}$: ACCELERATION
~~it~~ is the rate of change of velocity.

JERK: true! JERK is the rate of change of acceleration:
~~it's~~ an example of a third-derivative you can feel...

CURRENT: The CURRENT running through a point in a wire is $I(t) = \frac{dQ}{dt}$, where $Q(t)$ gives the flow of charge along the wire. inductance!


SLOPE: as you know from your prior exposure, a derivative is a slope ... **FALSE FALSE FALSE FALSE !!1!!**
The rate of change of y with respect to x is $\frac{dy}{dx}$, which has an interpretation as the tangent line slope.

CAVEAT! The word "RATE" is used in other contexts: unemployment rate is not a derivative, while marginal rate usually is!



OPERATOR

You should think of the derivative as a rate of change, or as part of a first-order approximation. But how should one interpret DIFFERENTIATION? The present setting is unique: $f: \mathbb{R} \rightarrow \mathbb{R}$ and $f': \mathbb{R} \rightarrow \mathbb{R}$ also. In pictures...



Differentiation is an OPERATOR, transforming a function f to its DIFFERENTIAL, df . The subject of differentials is full of devilish details. Suffice to say that you should think of df as the shorthand for the function that encodes the rate(s) of change of the output(s) of f , depending on the input(s) and rate(s) of change thereof. [see FLCT II for a better explanation of differentials...]

EXAMPLE: You know how to work with differentials, even if only via your prior exposure to integration: $u = \sin x \Rightarrow du = \cos x dx$




THE RULES of d

$d(f+g) = df + dg$	sum
$d(fg) = f dg + g df$	PRODUCT
$d\left(\frac{f}{g}\right) = \frac{g f' - f g'}{g^2}$	QUOTIENT
$d(f \circ g) = df \cdot dg$	CHAIN

The BKG IDEA is this: using differentiation as an OPERATOR allows you to transform equations by d into equations of rates. This is the basis for various RELATED RATES problems.

QUESTION: If round droplet of water evaporates at a rate proportional to its surface area. At the instant when the droplet vanishes, which is changing more rapidly, its surface area? its volume? its radius?



HINT: $V = \frac{4}{3}\pi r^3$; $A = 4\pi r^2$; $dA = C \cdot A$ keep going...

TRANSFORMS

Differentiation d , is not the only weapon to wield on equations: numerous other operators both respect equations and play nicely with differentiation.



One example is **LOGARITHMIC DIFFERENTIATION**, which exploits the **LOGARITHM** operator **LN** followed by **D**

EXAMPLE $y = x^x \xrightarrow{\text{LN}} \ln y = x \ln x \xrightarrow{\frac{d}{dx}} \frac{1}{y} dy = (\ln x) dx + x \left(\frac{1}{x} dx \right)$

$\frac{dy}{dx} = x^x (1 + \ln x) \xleftarrow{\text{ALGEBRA}} dy = y (1 + \ln x) dx$



Other examples demonstrate: **OPERATORS TRANSFORM EQUATIONS**

EXAMPLE $y = \lim_{x \rightarrow \infty} \left(1 + \frac{1}{x}\right)^x \xrightarrow{\text{LN}} \ln y = \lim_{x \rightarrow \infty} x \ln \left(1 + \frac{1}{x}\right) \quad [\text{as } x \rightarrow \infty]$


$\xrightarrow{\text{TAYLOR}}$ $\ln y = \lim_{x \rightarrow \infty} x \left(\frac{1}{x} - \frac{1}{2x^2} + \frac{1}{3x^3} + \text{HOT}\right)$

$\xleftarrow{\text{EXP}}$ $\ln y = 1$



This example shows how the operator **LN** allows for **Taylor expansion**. The inverse operator, **EXP**, recovers the solution.

this is a...
FORESHADOWING...
Transforms are important tools in control theory, signal processing, imaging & more...



YOU MAY FIND IT HELPS YOUR INTUITION TO APPLY WHAT YOU KNOW ABOUT DERIVATIVES: HERE FOLLOW SOME USES...

OPTIMIZATION I suspect you've seen basic optimization in the guise of **MAX-MIN** problems. These follow a common pattern:

STEP 1: DRAW A PICTURE, **STEP 2: DETERMINE ALL VARIABLES/CONSTRAINTS**



STEP 3: WRITE OUT THE FUNCTION TO BE EXTREMIZED


STEP 4: DIFFERENTIATE & FIND CRITICAL POINTS

STEP 5: CLASSIFY CRITICAL POINTS (max/min/?)

STEP: PROFIT.

The interesting part is classification...

To determine whether you have a max or a min, you will likely use the "SECOND-DERIVATIVE TEST", which is how high school students pronounce "TAYLOR SERIES". For $x=a$, a critical point of a function $f(x)$, one infers -



To grasp why this is true, Taylor expand @ $x=a$...

$$f(a+h) = \underbrace{f(a)}_{\text{VALUE}} + \cancel{f'(a)h} + \underbrace{\frac{1}{2}f''(a)h^2}_{\text{SIGN}} + \underbrace{\text{H.O.T.}}_{\text{QUADRATIC}}$$

(If $f'(a)$ is not defined, you must use other tools...)

This implies that near $x=a$, f is quadratic in $h = x-a$. If the second derivative also vanishes, you simply take the next term in the Taylor series (3rd, 4th, etc.)...


DIFFERENTIAL EQUATIONS

ODEs

An ORDINARY DIFFERENTIAL EQUATION on $x(t)$ is simply an algebraic equation implicating x and its derivative(s). These are singularly indispensable...



EXAMPLE | For $N(t)$ = number of bacteria in a nutrient-filled dish, one simple model for population dynamics is that N increases at a rate proportional to N . That is,



$$\frac{dN}{dt} = rN \quad (r > 0 \text{ RATE CONSTANT})$$


Let the reader note that $N(t) = Ce^{rt}$ is a solution for any C (constant), where $C = N(0)$. BUT... (1) This is an obviously flawed Malthusian model; (2) how do you solve this or more accurate ODE models?



~~STAY TUNED...~~

FLCT Chapter 3:

~~Integration~~



This chapter defines & computes integrals...

ANTI-DERIVATIVES The INDEFINITE INTEGRAL is easy to define: it is the inverse of the derivative.

$$\int f(x) dx := \{ F(x) \text{ such that } F'(x) = f(x) \} \quad +$$

Note: this is a class of functions, unique up to a constant. ↗

EXAMPLE

$$\int \frac{1}{(1-x)^2} dx = \frac{x}{1-x} + C$$

BUT

$$\int \frac{1}{(1-x)^2} dx = \frac{1}{1-x} + C$$

PROOF: If $F'(x) = G'(x)$ for all x , then
 $0 = F'(x) - G'(x) = (F-G)'(x)$
So that $F-G = \text{CONSTANT}$ (since $\frac{d}{dx} 0 = 0$) ✓

HINT: These C 's are not equal!



It is a remarkable fact that anti-derivatives exist: all (smooth) functions have an indefinite integral. However, it may not be in terms of simple functions!

FREE? COMPUTER?

SEPARATION Simple differential equations motivate

integrals best:



$$\frac{dx}{dt} = f(t) \quad \Rightarrow \quad x = \int f(t) dt$$

Given gravitational acceleration $a(t) = -g$
integrate for VELOCITY $v(t)$ and POSITION $x(t)$ via:
 $\frac{dv}{dt} = a \Rightarrow v(t) = -gt + v_0$ $\frac{dx}{dt} = v \Rightarrow x = -\frac{gt^2}{2} + v_0 t + x_0$

Slightly more complex models lead to these forms of ODE's:

$$\frac{dx}{dt} = f(x)$$

[AUTONOMOUS
1st ORDER ODE]

$$\frac{dx}{dt} = f(x) g(t)$$

[SEPARABLE
1st ORDER ODE]

These are ubiquitous...

EXAMPLES: WORDS

The amount of GREENGOO decays at a rate proportional to the amount present.

$$\frac{dg}{dt} = -Kg$$



The number of bunnies increases at a rate proportional to the population, and decreases at a rate proportional to its square.

$$\frac{dN}{dt} = rN - kN^2$$




Temperature changes at a rate proportional to the difference to the ambient temperature.

$$\frac{dT}{dt} = C(T_a - T)$$



Such equations are solved by separation & integration:

EXAMPLE: $\frac{dg}{dt} = -Kg$



SEPARATE
 $\frac{dg}{g} \neq dt$

$$\frac{dg}{g} = -Kdt$$

INTEGRATE
 $\int \frac{dg}{g} = \int -Kdt$

$$g = e^{-Kt+C}$$

VOLLA!
 $\ln g = -Kt + C$

$$e^{\ln g} = e^{-Kt+C}$$


$$g(t) = g_0 e^{-Kt}$$

This is what we found experimentally in Ch. 2!

COUPLED OSCILLATORS

Consider the case of two simple oscillators, modeled as an angle, say $\theta_1(t)$ and $\theta_2(t)$. Each represents a repetitive behavior: a metronome, or a vibrating string, or a beating heart. The simplest model of oscillation is:

$$\frac{d\theta_1}{dt} = \omega \quad \frac{d\theta_2}{dt} = \omega \quad [\omega = \text{FREQUENCY CONST}]$$



Solution: $\theta_1(t) = \omega t + C_1$; $\theta_2(t) = \omega t + C_2$

BUT: what if we couple them... put some small influence of θ_2 on θ_1 , and vice versa. SINUSOIDAL coupling is simplest:

$$\frac{d\theta_1}{dt} = \omega + \varepsilon \sin(\theta_2 - \theta_1)$$

$$\frac{d\theta_2}{dt} = \omega + \varepsilon \sin(\theta_1 - \theta_2)$$

{ if θ_2 is a bit ahead of θ_1 , it slows down θ_2 and speeds up θ_1 ... }

This COUPLED SYSTEM cannot be solved explicitly as above. But, we can analyze the PHASE $\phi(t) = \theta_2(t) - \theta_1(t)$ between the two:


$$\frac{d\phi}{dt} = \frac{d}{dt}(\theta_2 - \theta_1) = \frac{d\theta_2}{dt} - \frac{d\theta_1}{dt} = (\omega + \varepsilon \sin(-\phi)) - (\omega + \varepsilon \sin \phi) = -2\varepsilon \sin \phi$$

(LET'S TRY! $\frac{d\phi}{\sin \phi} = -2\varepsilon dt \Rightarrow \int \frac{d\phi}{\sin \phi} = -2\varepsilon t + C$ OR NOES! $\int \csc \phi d\phi = ?$)

This is not an easy integral. So... let's consider ϕ not too large... For ϕ near zero,


 $\frac{d\phi}{dt} \approx -2\varepsilon \phi$ LINEAR ODE $\phi \approx \phi_0 e^{-2\varepsilon t}$ THUS... $\phi \rightarrow 0$

This predicts that two coupled oscillators tend to **SYNCHRONIZE!** This is a remarkably observable result...




INTEGRATION TECHNIQUES

If you need to compute $\int f(x) dx$ and you don't recognize it, what do you do? There is no general algorithm, and even the computer might not help. There are some techniques worth knowing. They are:



} differentiation rules reversed
} fun with algebra




You should know that these exist, how to use them, and when each is useful. But this topic is not at the heart of Calculus.

u-SUBSTITUTION This is the Chain Rule run in reverse.

If you substitute $u = u(x)$ then $du = \frac{du}{dx} dx$ and

$$\int f(u) du = \int f(u(x)) \frac{du}{dx} dx$$

GIVEN change to



The difficulty is in choosing u well...

To be honest, your hand remembers this better than your head...

EXAMPLE: $\int \frac{(1 + \ln x)^3}{x} dx = \int u^3 du = \frac{1}{4} u^4 + C = \frac{1}{4} (1 + \ln x)^4 + C$

$u = (1 + \ln x) \Rightarrow du = \frac{1}{x} dx$



INTEGRATION BY PARTS This is a double-substitution method based on running the Product Rule in reverse.

PRODUCT RULE

$$d(uv) = u dv + v du$$

INTEGRATE

$$\int d(uv) = \int u dv + \int v du$$


$$\int d\Theta = \Theta$$

$$uv = \int u dv + \int v du$$

$$\int u dv = uv - \int v du$$

INTEGRATION BY PARTS FORMULA

LOVE



As with all substitution methods, the only difficulty lies in choosing u, v correctly...



EXAMPLE :

$$\begin{aligned}u &= x \\du &= dx \\dv &= e^x dx \\v &= e^x\end{aligned}$$

$$\int \underline{x} \underline{e^x dx} = \underline{\underline{x}} \underline{\underline{e^x}} - \int \underline{e^x} \underline{dx} = e^x(x-1) + C$$

EASY!

EXAMPLE : Let $I = \int e^x \sin x dx$. Computing I requires a double...

$$\begin{aligned}u &= \sin x \\du &= \cos x dx \\dv &= e^x dx \\v &= e^x\end{aligned}$$

$$\begin{aligned}u &= \cos x \\du &= -\sin x dx \\dv &= e^x dx \\v &= e^x\end{aligned}$$

$$I = e^x \sin x - \int e^x \cos x dx \quad J = \int e^x \cos x dx$$

$$J = e^x \cos x + \int e^x \sin x dx \quad I, \text{ again!}$$

DRASTIC! wait...

But combining these gives $I = e^x \sin x - (e^x \cos x + I)$ or

$$\begin{aligned}2I &= e^x(\sin x - \cos x) \\ \Rightarrow I &= \frac{1}{2} e^x(\sin x - \cos x) + C\end{aligned}$$



I DON'T

TRIGONOMICON There is a body of methods based on various trigonometric substitutions. You should be familiar with the use of these methods, but do not love them, for they will not love you back. They are merely tricks based on



$$\begin{aligned}1 - \sin^2 x &= \cos^2 x &\rightarrow u = \sin x & \left\{ \text{good for } \sqrt{1-x^2} \text{ terms} \right. \\ 1 + \tan^2 x &= \sec^2 x &\rightarrow u = \tan x & \left\{ \text{good for } \sqrt{1+x^2} \text{ terms} \right. \\ 1 + \sinh^2 x &= \cosh^2 x &\rightarrow u = \sinh x & \left. \begin{array}{l} \text{i told you he} \\ \text{was tricky...} \end{array} \right.\end{aligned}$$

EXAMPLE :

$$\int \frac{dx}{\sqrt{1+x^2}}$$

did I mention how much I love hyperbolic trigz?

Let $x = \tanh u$, $dx = \text{sech}^2 u du$

$$= \int \frac{\text{sech}^2 u du}{\sqrt{1+\tanh^2 u}} = \int \text{sech } u du$$


STUCK STUCK STUCK

Let $x = \sinh u$, $dx = \cosh u du$

$$= \int \frac{\cosh u du}{\sqrt{1+\sinh^2 u}} = \int du = u + C$$
 $= \text{ARCSINH } x + C$

PARTIAL FRACTIONS This, too, is an algebraic trick worth learning.

If you need to integrate a RATIONAL function (quotient of polynomials), and if the denominator factors nicely, you may split it up...



If $f(x) = \frac{p(x)}{(x-r_1)(x-r_2)\dots(x-r_n)}$ where the "roots" r_i are distinct and p is a polynomial of degree $< n$, then $f(x)$ splits up into a sum

$$f(x) = \frac{A_1}{x-r_1} + \frac{A_2}{x-r_2} + \dots + \frac{A_n}{x-r_n} \quad \text{for some constants } A_i$$

EXAMPLE : $f(x) = \frac{4x^2 - 3x - 4}{x^3 + x^2 - 2x}$ = $\frac{4x^2 - 3x - 4}{x(x-1)(x+2)}$ = $\frac{A}{x} + \frac{B}{x-1} + \frac{C}{x+2}$

put over a common denominator...

$$= \frac{A(x-1)(x+2) + Bx(x+2) + Cx(x-1)}{x(x-1)(x+2)} \stackrel{=}{\leftarrow} \frac{4x^2 - 3x - 4}{x(x-1)(x+2)} \stackrel{=}{\leftarrow}$$

$4x^2 - 3x - 4 = \underbrace{(A+B+C)x^2}_{4} + \underbrace{(A+2B-C)x}_{-3} + \underbrace{(-2A)}_{-4}$ 3 EQUATIONS,
3 UNKNOWN'S:
#WINNING

$$\left. \begin{array}{l} A = 2 \\ B = -1 \\ C = 3 \end{array} \right\} \Rightarrow \int \frac{4x^2 - 3x - 4}{x^3 + x^2 - 2x} dx = \int \frac{2}{x} - \frac{1}{x-1} + \frac{3}{x+2} dx = 2\ln|x| - \ln|x-1| + 3\ln|x+2| + C$$

NOTE: the algebra in solving for A, B, C is simpler if you substitute in specific values of $x = 0, 1, -2$. Try it!

STRATEGY & INTEGRATION Students want to know the secret algorithm or strategy for computing an integral in general. I will tell you.

BWAHAHAHAHAHAHAHAAH

If it does not exist.




You're on your own, kids.
Might want to try google/wiki/alpha.

DEFINITE INTEGRALS The indefinite integral is a (collection of) function (s, up to a constant), given by ~~anti~~ differentiation. This was motivated by finding solutions to differential equations. In contrast, the **DEFINITE INTEGRAL** is numerical and motivated by problems of geometry, physics, & statistics. You have no doubt computed lots of definite integrals in the past: it is now time to learn its meaning...

WARNING!
DON'T BE
FOOLED...



looks like the indefinite integral; however, it has an entirely different meaning!




DEFINITION: For an interval $[a, b]$, let \mathbb{P} denote a **PARTITION** into intervals $a = x_0 < x_1 < x_2 < \dots < x_{n-1} < x_n = b$. Let $\Delta x_i = x_i - x_{i-1}$, and choose, for each $i = 1 \dots N$ a **SAMPLE POINT** x_i^* in $[x_{i-1}, x_i]$. The **RIEMANN SUM** of f on \mathbb{P} is defined as:

$$\sum_{i=1}^n f(x_i^*) \Delta x_i$$

The **DEFINITE INTEGRAL** of f on $[a, b]$ is the limit of the Riemann sums over partitions \mathbb{P} on which $|\mathbb{P}| = \max(\Delta x_i) \rightarrow 0$.

$$\int_a^b f(x) dx = \lim_{|\mathbb{P}| \rightarrow 0} \sum_i f(x_i^*) \Delta x_i$$

unusual...
...interesting



A definite integral is therefore a very specific sort of limit-sum. You can see why it is often associated with an area; this interpretation is perhaps not the best (it will not survive multivariable calculus!). I prefer a **MASS ANALOGY**...

$f(x)$ = density of a rod at position x

$\int_a^b f(x) dx$ = mass of rod, $a \leq x \leq b$


THE FUNDAMENTAL THEOREM Although Riemann sums and limits thereof can be computed, it is a big pain, not unlike that of the computation of the derivative from the definition. It is much better to use the FTIC, which entwines the definite & indefinite:

THEOREM: Let $f(x)$ be continuous. Then ...

$$\textcircled{1} \quad \frac{d}{dx} \left(\int_a^x f(t) dt \right) = f(x); \text{ and}$$

$$\textcircled{2} \quad \int_a^b f(x) dx = \left[\int f(x) dx \right]_a^b \quad \begin{matrix} \text{evaluated from} \\ a \text{ to } b \dots \end{matrix}$$

DEFINITE INTERVAL INDEFINITE INTERVAL



Proof: $\textcircled{1} \quad \frac{d}{dx} \left(\int_a^x f(t) dt \right) = \lim_{h \rightarrow 0} \frac{1}{h} \left(\int_a^{x+h} f(t) dt - \int_a^x f(t) dt \right) = \lim_{h \rightarrow 0} \frac{1}{h} \int_x^{x+h} f(t) dt = \lim_{h \rightarrow 0} \frac{1}{h} f(x^*) \Delta x = f(x)$

$\textcircled{2}$ Since $\int f(t) dt$ is an antiderivative of $f(x)$, and since all anti-derivatives differ by a constant, then, for $G(x)$ any antiderivative of f ,

$$\int_a^x f(t) dt - G(x) = C \quad \text{CONSTANT}$$

evaluate $\int_a^x f(t) dt - G(a) = C$ $\xrightarrow{\text{thus}} C = -G(a)$ and
by definition



$$\begin{aligned} \int_a^b f(t) dt &= G(b) + C \quad \text{for any antiderivative } G \\ &= G(b) - G(a) \quad " " \\ &= \left[\int f dx \right]_a^b \end{aligned}$$

Quad-Erat-Demonstration
= "for thizle" in Latin

This sets the stage for using this chapter's tools for definite integrals ...

EXAMPLE


$$\begin{aligned} \int_{x=a}^b f(u(x)) \frac{du}{dx} dx &= \int_{u=a}^{u(b)} f(u) du \quad u\text{-SUBS} \\ \int_{x=a}^b u(x) dv(x) &= u(x)v(x) \Big|_{x=a}^b - \int_{x=a}^b v(x) du(x) \quad \text{PARTS} \end{aligned}$$


Next up...

WHAT'S IT ALL GOOD FOR?

FLCT Chapter 4:

~~Applications~~





...and all the Calculus learned thus far is good for...?



This Chapter surveys some typical applications...

THE BIG IDEA

it's very simple...

$$\text{Desideratum} = \int \text{Rate} d\text{Desideratum}$$

Desideratum = Desideratum

the "Desideratum"-element


If you want something, determine its ELEMENT -- its rate of change -- and integrate.

MASS Consider a rod of varying density $\rho(x)$

WHAT IS ITS MASS, m ?

MASS ELEMENT

$$\text{MASS} \quad m = \int_{x=a}^b dm = \int_{x=a}^b \rho(x) dx$$



"So, I can think of dm as an itty bitty mass, right?"

WELL... NO. IT'S A DIFFERENTIAL. BUT, WHATEV.

LENGTH


What is the length of a curve?

If you have a parametrized curve...

$$\gamma(t) = (x(t), y(t))$$

$$dl = \sqrt{dx^2 + dy^2}$$

$$= \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} dt$$



If you have a graph...

$$y = f(x)$$


$$dl = \sqrt{dx^2 + dy^2}$$

$$= \sqrt{1 + \left(\frac{dy}{dx}\right)^2} dx$$

$$l = \int dl$$

AREA

Here is a simple example of the principle: AREA OF A DISC (radius r)




$$dA = \frac{1}{2}(r d\theta)r$$

$$= \frac{1}{2}r^2 d\theta$$

$$A = \int dA = \int_{\theta=0}^{2\pi} \frac{1}{2}r^2 d\theta = \pi r^2$$

π

This is how Archimedes did it. kinda.




$$dl = 2\pi x dx$$

$$A = \int dl = \int_{x=0}^r 2\pi x dx = \pi r^2$$



...or, use a different area element



Yes, your favorite example, the area between two curves, works:

$$A = \int dA = \int_{x=a}^b (f(x) - g(x)) dx$$

DON'T FORGET! Sometimes using horizontal instead of vertical is better!


SURFACE AREA

The area of a non-flat surface takes some effort to compute. For simplicity, consider a surface obtained by revolving a curve about an axis: a **SURFACE OF REVOLUTION**...

$$ds = \sqrt{r^2 + (dr/dx)^2} dx$$

SURFACE AREA ELEMENT DISTANCE TO AXIS ARCLENGTH ELEMENT

$$S = \int ds$$




If you rotate $y = f(x)$ about the x -axis, then $r = f(x)$ and $dl = \sqrt{1 + \left(\frac{dy}{dx}\right)^2} dx$

VOLUME As you will have guessed by now ...
and the challenge is to find a good VOLUME ELEMENT

$$V = \int dV$$

what is that symbol?
sign you keep writing
in your notes?

You may recall the volume of a CONE over a disc of radius r and height h is $V = \frac{1}{3}\pi r^2 h$.



Why THE $\frac{1}{3}$? The slicing method gives

$$dV = A(y)dy = \pi x^2 dy = \pi r^2 \left(1 - \frac{y}{h}\right)^2 dy \quad \text{so...}$$



$$V = \int_{y=0}^h \pi r^2 \left(1 - \frac{y}{h}\right)^2 dy = \pi r^2 \int_{u=0}^1 h u^2 du = \pi r^2 h \left[\frac{u^3}{3}\right]_0^1 = \frac{1}{3}\pi r^2 h$$

But... this generalizes greatly to the volume of a cone of height h over an area B

AH! The $\frac{1}{3}$ came from πr^2

$$V = \frac{1}{3}Bh$$

CAN YOU DO THIS?



Now for an "improper" example...

ROTATE
 $y = x^p$
ABOUT
 x -**AXIS**
 $1 \leq x < \infty$

VOLUME


$$V = \int_{x=1}^{\infty} \pi \left(\frac{1}{x^p}\right)^2 dx = \begin{cases} \frac{\pi}{1-2p} x^{1-2p} \Big|_1^{\infty} & (p \neq \frac{1}{2}) \\ \pi \ln x \Big|_1^{\infty} & (p = \frac{1}{2}) \end{cases} = \begin{cases} \frac{\pi}{2p-1} & (p > \frac{1}{2}) \\ \infty & (0 < p < \frac{1}{2}) \\ \infty & (p = \frac{1}{2}) \end{cases}$$

SURFACE AREA

$$y = \frac{1}{x^p}$$

$$S = \int_{x=1}^{\infty} 2\pi x \sqrt{1 + \left(\frac{p}{x^{p+1}}\right)^2} dx = ? \quad \text{ESTIMATE!}$$

$$\geq \int_{x=1}^{\infty} 2\pi x \sqrt{\left(\frac{p}{x^{p+1}}\right)^2} dx = \int_{x=1}^{\infty} \frac{2\pi p}{x^p} dx = \begin{cases} \frac{2\pi p}{p-1} & p > 1 \\ \infty & p \leq 1 \end{cases}$$




BALLS

Let's compute the "volume" of an n -dimensional ball, radius r

WHAT IS... THE n^{th} DIMENSION?

$\mathbb{R}^n = \{(x_1, \dots, x_n) \text{ with each } x_i \in \mathbb{R}\}$
ordered n -tuples of reals...
The n -dimensional ball is
 $B^n(r) = \{(x_1, \dots, x_n) : x_1^2 + \dots + x_n^2 \leq r^2\}$



IDEA: Use the "slicing" method
A "hyperplane" cuts the ball into
a ball of one dimension less. As
you sweep across, the radius of the
($n-1$)-dimensional ball equals

$$r = \sqrt{1-x^2} \quad -1 \leq x \leq 1$$

(assuming the original is a unit n -ball)



Let $V_n(r) = \text{"volume"} \text{ of } n\text{-dimensional ball of radius } r$, with $V_n := V_n(1)$

LEMMA: $V_n(r) = V_n r^n$ (think: units \notin scaling) $\Rightarrow =: C_n \text{ CONSTANT}$

$$\text{LEMMA: } V_n = \int_{x=-1}^1 V_{n-1}(\sqrt{1-x^2})^{n-1} dx = V_{n-1} \int_{x=-1}^1 (1-x^2)^{\frac{n-1}{2}} dx$$

$$V_n = C_n V_{n-1}$$

Using the substitution $x = \sin \theta$ and integration by parts, one obtains ...

$$C_n = \int_{\theta=0}^{\pi} (1-\sin^2 \theta)^{\frac{n-1}{2}} \cos \theta d\theta = \int_{\theta=0}^{\pi} \cos^n \theta d\theta = \frac{n-1}{n} \int_{\theta=0}^{\pi} \cos^{n-2} \theta d\theta$$

i.e.,

$$C_n = \frac{n-1}{n} C_{n-2}$$

By hand, one computes ...

This gives an inductive approach for computing C_n
with some effort, an induction argument yields...

The volume of the
unit ball in dimension
 n equals...

$$V_n = \begin{cases} \frac{\pi^k}{k!} & : n=2k \text{ EVEN} \\ \frac{2\pi^k k!}{n!} & : n=2k+1 \text{ ODD} \end{cases}$$

QUESTION What is $\lim_{n \rightarrow \infty} V_n$? Are you frightened?

n	C_n	V_n
0	π	1
1	2	2
2	$\frac{\pi^2}{2}$	π^2
3	$\frac{4}{3}\pi$	$\frac{4}{3}\pi$
4	$\frac{8\pi^2}{8}$	$\frac{\pi^2}{2}$
5	$\frac{16}{15}\pi^2$	$\frac{16}{15}\pi^2$
6	$\frac{5\pi^3}{16}$	$\frac{\pi^3}{6}$
7	$\frac{32}{35}\pi^3$	$\frac{16}{105}\pi^3$
8	$\frac{35\pi^4}{128}$	$\frac{\pi^4}{24}$

LOSSY... AREA... VOLUME... HYPERVOLUME...

AVERAGES One of the best applications of integrals is in defining the average, or **MEAN**, of a function. The mean of $f(x)$ over $a \leq x \leq b$ is defined to be

$$\bar{f} = \frac{\int_a^b f(x) dx}{\int_a^b dx}$$

why write the denominator thus?
BIG DUH $\int_a^b dx = b-a$

It makes more sense in a little while...

PLEASE STAND BY


EXAMPLE: The **ROOT MEAN SQUARE** of f is $RMS(f) = (\bar{f^2})^{1/2}$

DUE TO POWER... $RMS(A \sin \omega t) = \left(\int_0^{2\pi/\omega} (A \sin \omega t)^2 dt \right)^{1/2} = \sqrt{A^2 - \frac{A^2}{2}} = \frac{A}{\sqrt{2}}$

CENTERS What is the exact center of a given body? It is the **CENTROID**, the point whose coordinates are averages. Consider the region $R = \{(x, y) : a \leq x \leq b, g(x) \leq y \leq f(x)\}$ between two curves. Its centroid is the point (\bar{x}, \bar{y}) given by the average x and y values over R .

THIS CALLS FOR TWO INTEGRALS, IN SUCCESSION:

"MAKE MAKE A DOUBLE"




$$\begin{aligned}\bar{x} &= \frac{\int_R x dA}{\int_R dA} = \frac{\int_{x=a}^b \left(\int_{y=g(x)}^{f(x)} x dy \right) dx}{\int_{x=a}^b dA} = \frac{\int_{x=a}^b x (f(x) - g(x)) dx}{\int_{x=a}^b f(x) - g(x) dx} \\ \bar{y} &= \frac{\int_R y dA}{\int_R dA} = \frac{\int_{x=a}^b \left(\int_{y=g(x)}^{f(x)} y dy \right) dx}{\int_{x=a}^b dA} = \frac{\int_{x=a}^b \frac{1}{2} [(f(x))^2 - (g(x))^2] dx}{\int_{x=a}^b f(x) - g(x) dx}\end{aligned}$$

These formulae give **CENTROIDS**: to get **CENTER-OF-MASS**, use a **DENSITY** $\rho(x, y)$ and do all integrals with respect to the **MASS ELEMENT** $dm = \rho dA$


MOMENTS

For an object rotated about an axis, the RMS of distance to the axis is called **RADIUS OF GYRATION**. The numerator



$I = \int r^2 dm$ is called the **MOMENT OF INERTIA**.

This moment represents the resistance to rotation about the axis...



TICKLE ME CTHULHU

PROBABILITY

A simple approach to probability is via MEASURES

COUNTING	Odds that a fair die lands on an odd number? $\frac{\text{ODDS}}{\text{TOTAL}} = \frac{3}{6} = \frac{1}{2}$
----------	--


LENGTH	Odds that a random angle has $\sin \geq \frac{1}{2}$? $\frac{\pi/3}{2\pi} = \frac{1}{3}$
--------	--

AREA	Odds that a random point in a square lies in the inscribed circle? $\frac{\pi r^2}{(2r)^2} = \frac{\pi}{4}$
------	--

VOLUME	Hyper-volume is the volume of a random variable in n-space
--------	---

Let's try it out...

THE BUNNEN NEEDLE PROBLEM




Given a needle of length 1, drop it on notebook paper with unit-distance lines. With what probability does the needle cross one of the lines?

The natural variables are: h , the distance from the low tip to the next line up; and θ , the angle.

NOTE: $0 \leq h < 1$ & $0 \leq \theta < \pi$


These variables are UNIFORM and INDEPENDENT: one has a crossing iff



what are the odds of a random (h, θ) satisfying this?

$$\frac{\int_0^\pi \sin \theta d\theta}{\pi \cdot 1} = \frac{2}{\pi}$$

THE AREA FRACTION



PROBABILITY THAT A "RANDOM" x LIES BETWEEN a AND b

Typical random variables are not uniform, but vary according to a PDF: PROBABILITY DENSITY FUNCTION, $p(x)$, satisfying:

$$P\{a \leq x \leq b\} = \int_{x=a}^b p(x) dx$$

Of course, a PDF must satisfy

$$\int_{-\infty}^{\infty} p(x) dx = 1$$

PROBABILITY THAT $-\infty < x < \infty$

Many definitions in probability & statistics are expressed in terms of integrals:

MEAN

$$\mu = \int x p(x) dx$$

The "average" value of x

HEY! THAT LOOKS LIKE CENTER-OF-MASS!

VARIANCE

$$V = \int (x - \mu)^2 p(x) dx$$

How "spread out" the distribution is

HEY! THAT IS THE MOMENT OF INERTIA ABOUT THE μ -axis!

STANDARD DEVIATION

$$\sigma = \sqrt{V}$$

Like V , but with better units

HEY! THAT'S THE RADIUS OF GYRATION ABOUT $x = \mu$!

Thinking of $p(x)dx$ as a

MASS ELEMENT dm has some advantages...

but that's not normal!

WORK As you know, work equals FORCE · DISTANCE; but this is only true locally; one needs to integrate the **WORK ELEMENT** in general.

SPRING PULLING...



If the force required to displace a spring from equilibrium ($x = 0$) is $F(x)$, then the work element is...

$$dW = F(x) dx$$




ROPE LIFTING...

To pull an object whose weight $w(y)$ varies with height y ... $dW = w(y) dy$

FLUID PUMPING...

To pump a fluid of density ρ , the work element is...



$$dW = h(z) A(z) \rho dz$$

net height pumped

cross-sectional area

LEARN SOMETHIN'


FORCE

A fluid exerts a **FORCE** on a submerged surface equal to **PRESSURE** times area, orthogonal to the surface. The pressure (and thus force) varies with depth and equals ρz , where z is depth and ρ is density. The force element is...



$$dF = \rho z dA$$

area element



DIFFERENTIAL EQUATIONS

We initially motivated indefinite integrals via differential equations, but definite integrals are useful, too, for **INITIAL VALUE PROBLEMS**: in the separable setting, one gets...

GIVEN: $\frac{dy}{dt} = f(y)g(t)$ THE DIFFERENTIAL EQUATION

$y(t_0) = y_0$ THE INITIAL CONDITION t_1 THE FINAL TIME


FIND: $y_1 := y(t_1)$

SOLUTION: Integrate and solve...

FOR SEPARABLE ODE'S THE REQUISITE 'ELEMENT' IS GIVEN TO YOU A PRIORI!


$$\int_{y=y_0}^{y_1} \frac{dy}{f(y)} = \int_{t=t_0}^{t_1} g(t) dt$$


THE MORAL Everybody wants to have a black box formula for solving stuff: do not do that. Construct an "element" and integrate. Carefully.



FLCT Chapter 5:

~~Discretization~~





Calculus is built for continuous objects; the world is full of discrete, quantized things. The synthesis of smooth & discrete Calculus is the focus of this final chapter of FLCT...

SEQUENCES

In single-variable calculus, the typical function has one real input and one real output: $f: \mathbb{R} \rightarrow \mathbb{R}$




The most common discretization is from \mathbb{R} to \mathbb{N} ,


the "natural" numbers, $\mathbb{N} = \{0, 1, 2, 3, 4, \dots\}$. The discrete analogue of a function is a **SEQUENCE**, $a: \mathbb{N} \rightarrow \mathbb{R}$, whose input is discrete. Thus,

$$a = (a_0, a_1, a_2, a_3, \dots)$$

$a(0) \uparrow \quad a(1) \uparrow \quad \text{etc.}$



EXAMPLE: Unemployment rate as a function of month




Exam scores as a function of student

Dow Jones index as a function of day

LIMITS


The first step in Calculus is **limits**. Relax... Σ

There are no tricky definitions here. One says $\lim_{n \rightarrow \infty} a_n = L$ iff a_n gets as close as you like to L for n sufficiently large.



The same deal holds as in other limit problems:

LEMMA: If $a = (a_n)$ is **BOUNDED** and **MONOTONE**, then $\lim_{n \rightarrow \infty} a_n$ exists!




EXISTENCE MUST BE ESTABLISHED!



Nope.


EXAMPLE

$$\sqrt{1 + \sqrt{1 + \sqrt{1 + \sqrt{1 + \sqrt{1 + \sqrt{\dots}}}}}} = \frac{1 + \sqrt{5}}{2} = 1 + \frac{1}{1 + \frac{1}{1 + \frac{1}{\dots}}}$$



$$\begin{aligned} a_{n+1} &= \sqrt{1 + a_n} && \text{RECURSION RELATION} & a_{n+1} &= 1 + \frac{1}{a_n} \\ L &= \sqrt{1 + L} && \text{LET } L = \lim_{n \rightarrow \infty} a_n & L &= 1 + \frac{1}{L} \\ L^2 &= 1 + L && \text{SOLVE FOR } L > 0 & L^2 &= L + 1 \end{aligned}$$

These sequences a_n are bounded and increasing... L exists!



Fortunately, one may use continuous techniques to establish limits in discrete settings...

EXAMPLE

$$\lim_{n \rightarrow \infty} n \sin \frac{1}{n} = \lim_{n \rightarrow \infty} \frac{\sin \frac{1}{n}}{\frac{1}{n}} = \lim_{x \rightarrow 0^+} \frac{\sin x}{x}$$

NOTE: $x = \frac{1}{n}$ are in \mathbb{R} , not \mathbb{N} ...

STANDARD	TAYLOR	L'HOPITAL	STUFF
----------	--------	-----------	-------

in so lonely...


Your knowledge of Taylor series & L'Hopital's rule is fair game...

(Is it strange that one can use derivatives to compute limits of sequences?
(Even though you cannot differentiate with respect to n ? Well...)

DERIVATIVES

What is the analogue of the derivative for sequences?

These are called FORWARD/BACKWARD DIFFERENCES



$$\Delta a_n = a_{n+1} - a_n$$

$$\nabla a_n = a_n - a_{n-1}$$

FORWARD
BACKWARD

Notice how Δa and ∇a are sequences that express rates of change as you increase (Δ) or decrease (∇) the input by 1.
(I wonder what happens if you average the two? Hmmm...)

ΨΜ
ΨΛΨ ΙΠ ΞΠ ΛΨ
ΦΙΨΙΨΗΦΙΝ...

There is an entire calculus for difference operators; Δ, ∇
(e.g.) If (a_n) and (b_n) are sequences, then there is a "product rule" for differences of $(ab)_n := (a_n b_n)$

$$\Delta(ab)_n = a_n \Delta b_n + b_n \Delta a_n + \Delta a_n \Delta b_n$$

$$\Delta(ab)_n = a_n \Delta b_n + b_n \Delta a_n - \Delta a_n \Delta b_n$$

This, as you might guess, is just the beginning...

IN PRINCPIO


DIFFERENTIAL EQUATIONS if you can define "derivatives" of a sequence, why not differential equations?

$$\frac{dx}{dt} = f(x) \quad \text{cf.} \quad \Delta x_n = f(x_n)$$

DIFFERENTIAL
vs.
DIFERENCE

$$\approx x_{n+1} = x_n + f(x_n)$$

which, since
 $\Delta x_n = x_{n+1} - x_n$,



Any "initial condition" x_0 yields a discrete-time solution sequence (x_n) via iteration

These "difference equations" are very useful in continuous settings:

NEWTON'S METHOD


GIVEN: $f: \mathbb{R} \rightarrow \mathbb{R}$ continuous

PROBLEM: find a root of f

CHOOSE: starting guess x_0

SOLUTION: iterate the equation

$$x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)}$$



INTUITION: cf. $\frac{dx}{dt} = -\frac{f(x)}{f'(x)}$

CAVEAT: You may ALWAYS MOVE TOWARD ROOT not converge to the root you wanted, nor to anything!

EXAMPLE: $f(x) = 1 - x^2$

$$\text{if } x_0 \text{ is } \begin{cases} < 0 : x_n \rightarrow -1 \\ = 0 : x_1 = \#FAIL! \\ > 0 : x_n \rightarrow +1 \end{cases}$$

EULER'S METHOD

GIVEN: ODE $\frac{dx}{dt} = f(x, t)$

initial condition $x_0 = x(t_0)$

PROBLEM: find $x(T)$ for $T > t_0$

CHOOSE: step size st ($= T - t_0 / N$)

SOLUTION: iterate the equation

$$x_{n+1} = x_n + f(x_n, t_n) \\ t_{n+1} = t_n + st$$

INTUITION: cf. $\frac{dx}{dt} = f(x, t)$
TIME KEEPS ON (SLIPPING)³ INTO THE FUTURE $\frac{dt}{dt} = 1$

CAVEAT: When you get to T , you have only an approximation to $x(T)$, the accuracy of which depends upon st .

EXAMPLE: $\frac{dx}{dt} = x \quad x_0 = 1 \quad t_0 = 0$

We know $x(t) = e^t$ so $x(1) = e$.

CHOOSE: $st = 1/N$; then

$$x_{n+1} = x_n + x_n/N = x_n(1 + 1/N)$$

$$t_{n+1} = t_n + 1/N = n/N$$

$x_N = (1 + 1/N)^N$ approximates e

As $st \rightarrow 0$, $N \rightarrow \infty$ and

$$\lim_{N \rightarrow \infty} x_N = \lim_{N \rightarrow \infty} (1 + \frac{1}{N})^N = e$$

"IS THAT ALL THERE IS?" NOPE. TAKE A NUMERICAL ANALYSIS COURSE.

INTEGRALS


The analogue of the definite integral for a (finite) sequence $\{a_n\}$ is simply the sum $\sum_{n=0}^k a_n$. There's nothing fishy here, unless your goal is to do

NUMERICAL INTEGRATION -- the approximation of $\int_a^b f(x) dx$ from a discrete sampling of f ... this requires some choices!


CHOOSE: step size $\Delta x = (b-a)/N$ for some N

DEFINE: sample points $x_n = a + n \cdot \Delta x$ for $n = 0 \dots N$

You can approximate the integral by using the x_i 's to sample f :



These give simple-to-compute, but coarse, approximations. To better approximate, use a Taylor series approach, approximating f locally with polynomials



SERIES The analogue of an improper integral ($\int_0^\infty f(x)dx$) is:

$$\sum_{k=0}^{\infty} a_k = a_0 + a_1 + a_2 + \dots$$

INTEGRAL TEST

BEATIFIC! $1 + \frac{1}{2} + \frac{1}{4} + \frac{1}{8} + \frac{1}{16} + \dots = 2$

ENIGMATIC! $1 - 1 + 1 - 1 + 1 - 1 + \dots = ?$

is it zero? one?
or $\frac{1}{1-e} = \frac{1}{e}$?

What to do? Use limits, as with the rest of Calculus.

DEFINE: $S_n = \sum_{k=0}^n a_k = a_0 + a_1 + \dots + a_n \quad \left\{ \begin{array}{l} \text{"PARTIAL SUMS"} \\ \text{FULL PATHONIC PATH: MY SISTER LIVES THERE ARE PEOPLE THAT WORK IN IT"} \end{array} \right.$

$$\sum_{k=0}^{\infty} a_k := \lim_{n \rightarrow \infty} S_n \quad \left\{ \begin{array}{l} \text{"CONVERGES" or} \\ \text{"DIVERGES"} \end{array} \right.$$

This is the proper definition, and fits with our notion of the **SERIES** as a discrete integral...

$$\int_0^\infty f(x)dx = \lim_{T \rightarrow \infty} \int_0^T f(x)dx$$

\uparrow Cf. \downarrow

$$\sum_{k=0}^{\infty} a_k = \lim_{L \rightarrow \infty} \sum_{k=0}^L a_k$$



QUESTIONS:

- 1: Does a given series $\sum_{k=0}^{\infty} a_k$ converge?] NOT SO HARD..
- 2: If so, to which value does it converge?] NOT SO EASY...
- 3: Why is an infinite sum called a "series"?] I'LL TELL YOU LATER..

CONVERGENCE TESTS

There are numerous tests for series to ascertain convergence or divergence. Be careful of hypotheses!

DIVERGENCE TEST

HYPOTHESES: none

TEST: if $\lim_{n \rightarrow \infty} a_n \neq 0$, then $\sum_n a_n$

DIVERGES

EXAMPLE: $\sum_{n=1}^{\infty} \cos(n\pi)$ DIVERGES!
 $= 1 - 1 + 1 - 1 + 1 - 1 + \dots$

COMPARISON TEST

HYPOTHESES: $a_n \geq 0$; $b_n \geq 0$ TEST:

if $a_n \leq b_n$ and $\sum_n b_n$ CONVERGES DIVERGES then

$b_n \leq a_n$ also CONVERGES DIVERGES since $\sum_n a_n \leq \sum_n b_n$
 $\sum_n b_n \leq \sum_n a_n$

EXAMPLE:

$$\sum_{n=1}^{\infty} \frac{\ln(n)}{n} \stackrel{\text{DIV}}{\geq} \sum_{n=1}^{\infty} \frac{1}{n} = \infty \quad \sum_{n=1}^{\infty} \frac{\sqrt{n-1}}{n^2} \stackrel{\text{CONV}}{\leq} \sum_{n=1}^{\infty} \frac{1}{n^{3/2}} < \infty$$


INTEGRAL TEST

HYPOTHESES: $a_n \geq 0$, decreasing

TEST: choose $a(x)$ continuous, decreasing, with $a(n) = a_n$. Then

$$\sum_n a_n \stackrel{\text{CONVERGE OR DIVERGE TOGETHER!}}{\longrightarrow} \int_1^{\infty} a(x)dx$$

EXAMPLE:





LIMIT TEST	RATIO TEST	ROOT TEST
<p><u>HYPOTHESES:</u> $a_n \geq 0, b_n \geq 0$</p> <p><u>TEST:</u> if $0 < \lim_{n \rightarrow \infty} \frac{a_n}{b_n} < \infty$</p> <p>then $\sum a_n$ CONVERGE OR DIVERGE TOGETHER! $\sum b_n$</p> <p><u>EXAMPLE:</u> $\sum \frac{n^2 - 2n + 3}{5^{n-1} + n^3}$ CONVERGES since $\sum \left(\frac{3}{5}\right)^n$ does...</p>	<p><u>EXAMPLE:</u> $\sum \frac{n!}{1 \cdot 3 \cdot 5 \cdots (2n-1)}$</p> <p>DIVERGES since $P = \frac{1}{2}$</p> <p><u>EXTRA:</u> $\sum \frac{1}{n^p}$ DIVERGES if $P \leq 1$ CONVERGES if $P > 1$</p>	<p><u>HYPOTHESES:</u> Compute some L!</p> <p><u>TEST:</u> $L = \lim_{n \rightarrow \infty} \sqrt[n]{ a_n }$</p>


It takes a lot of practice to know which test to use when...
Some types of series are not so bad, however...

ALTERNATING SERIES

An alternating series alternates $\sum_{n=0}^{\infty} (-1)^n a_n$ (with $a_n \geq 0$) positive and negative terms. For such, it is easy to determine convergence: An alternating series (with decreasing terms a_n) converges IF & ONLY IF the limit $\lim_{n \rightarrow \infty} a_n = 0$



PROOF:
Use the number line bunny



Partial sums hop back & forth by length a_n

CONVERGENCE

There are degrees of convergence; one says
 $\sum a_n$ CONVERGES ABSOLUTELY if $\sum |a_n|$ CONVERGES. A series
 $\sum a_n$ CONVERGES CONDITIONALLY if it converges but not absolutely.

EXAMPLES: $\sum_n \frac{(-1)^n}{n^p}$

DIVERGES
CONVERGES CONDITIONALLY
CONVERGES ABSOLUTELY

$p \leq 0$
 $0 < p \leq 1$
 $1 < p$


divergence test
alternating test
 p -series test



POWER SERIES

The theme of this chapter is the discretization of functions to sequences and the resulting Calculus. It is time to reverse the process. A **POWER SERIES** in x is

$$\sum_{n=0}^{\infty} a_n x^n$$




for some coefficients a_n . This converts a series into a function of x (at least when the series converges!). When does it converge?



$$\text{Let } p = \lim_{n \rightarrow \infty} \left| \frac{a_{n+1} x^{n+1}}{a_n x^n} \right| = |x| \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right|$$

Then $f(x) = \sum_n a_n x^n$ CONVERGES ABSOLUTELY if $p < 1$ (and DIVERGES if $p > 1$) by the ratio test.

EST Let $R = \lim_{n \rightarrow \infty} \left| \frac{a_n}{a_{n+1}} \right|$. Solving $p < 1$ yields $|x| < R$; hence, R is called the **RADIUS OF CONVERGENCE**. So, a power series behaves like so,



R can be any number, such as:

- ∞ for $a_n = \frac{1}{n!}$
- 1 for $a_n = 1$
- 0 for $a_n = n!$

The Taylor series of Chapter 1 return. For $f(x) = \sum_n a_n x^n$ and $|x| < R$, the Taylor series of f is, precisely, the power series...

BUT: you must be careful at $x = \pm R$!

Power series of the form $\sum_n a_n (x-c)^n$ have an INTERVAL OF CONVERGENCE centered at c

EXAMPLE: By taking $\lim_{x \rightarrow 1^-} \ln(1+x)$, we get...

$$\text{RECALL: for } |x| < 1 \quad \ln(1+x) = \sum_{n=1}^{\infty} (-1)^{n-1} \frac{x^n}{n}$$

DIVERGES AT $x = -1$
CONVERGES AT $x = 1$

$$\ln 2 = 1 - \frac{1}{2} + \frac{1}{3} - \frac{1}{4} + \frac{1}{5} - \frac{1}{6} + \frac{1}{7} - \frac{1}{8} + \frac{1}{9} - \frac{1}{10} + \dots$$

$$\frac{1}{2} \ln 2 = \frac{1}{2} - \frac{1}{4} + \frac{1}{6} - \frac{1}{8} + \frac{1}{10} - \frac{1}{12} + \dots$$

$$\frac{3}{2} \ln 2 = 1 + \frac{1}{3} - \frac{1}{2} + \frac{1}{5} + \frac{1}{7} - \frac{1}{4} + \frac{1}{9} + \frac{1}{11} - \frac{1}{6} + \dots$$

Nooooooo....! These are the same terms, just in a different order...



Yes, dear reader, a conditionally convergent series can be rearranged to sum up to ANY NUMBER YOU WANT. Beware! Fortunately, absolutely convergent series behave as nicely as can be...

TAYLOR, AGAIN All this work, just to come back to Taylor series?
 Yep. But notice the language we have developed; ex gratia-

THEOREM: Within the radius of convergence of a power series for $f(x)$, one can differentiate and integrate f or the series equivalently.

This, then, is a synthesis of discrete and continuous Calculus: one can evaluate a discrete "integral" by taking a Taylor series and using all the tools & tricks of Calculus:

EXAMPLE: Evaluate the geometric series at $-x^2$ to obtain

$$\frac{1}{1+x^2} = 1 - x^2 + x^4 - x^6 + x^8 - \dots \quad |x|<1$$

Integration yields:


$$\text{ARCTAN}(x) = x - \frac{x^3}{3} + \frac{x^5}{5} - \frac{x^7}{7} + \dots \quad |x|<1$$

Taking the limit as $x \rightarrow 1^-$ yields...


$$1 - \frac{1}{3} + \frac{1}{5} - \frac{1}{7} + \dots = \frac{\pi}{4} \quad \text{SURPRISE!}$$



THE END? The use of power series relies on polynomials as a basis for approximation. Different bases are possible:



FOURIER SERIES



WAVELETS

The most exciting thing in Calculus is how the same basic ideas recur, no matter how deep one treads. This will be evident when you explore multivariable Calculus...

These fragments I have
shoved against my ruins...

