

Math 334 Sample Problems

One side of one notebook sized page of notes will be allowed on the test. You may work together on the sample problems – I encourage you to do that. The test will cover up to §3.4.

1. Prove that the level set $y^2 = (x - 1)(x - 2)(x - 3)$ is the disjoint union of two smooth connected curves. Prove that one of the curves is compact and the other is not.
2. Suppose that (x_0, y_0, z_0, u_0) satisfies the equations

$$\begin{aligned}x + y + z &= F(u) \\x^2 + y^2 + z^2 &= G(u) \\x^3 + y^3 + z^3 &= H(u),\end{aligned}$$

where, F, G, H are C^1 in a neighborhood of u_0 . State a sufficient condition for being able to solve these equations for x, y, z as C^1 functions of u in a neighborhood of (x_0, y_0, z_0, u_0) .

3. Is the set $\{(x, y) : y^2 + x^2 e^y = 0\}$ a smooth curve? Is the set $\{(a \cos t, b \sin t) : t \in (0, \pi)\}$, where $a > 0, b > 0$ a smooth curve?
4. Expand $(1 - x + 2y)^3$ in powers of $x - 1$ and $y - 2$ in two different ways. The first way is by using algebra and the second way is by computing the Taylor polynomial of degree three centered at $(1, 2)$.
5. Using the method of Lagrange multipliers, find the highest and lowest points of the circle

$$x^2 + y^2 + z^2 = 16, \quad (x + 1)^2 + (y + 1)^2 + (z + 1)^2 = 27$$

6. Show that the surface $z = 3x^2 - 2xy + 2y^2$ lies entirely above every one of its tangent planes. Hint: Look at the Taylor expansion at every point.
7. Let $a > 0, b > 0$ and $a + b = 1$. Also let $x > 0, y > 0$. prove that

$$x^a y^b \leq ax + by,$$

by using the method of Lagrange multipliers applied to maximize $x^a y^b$ subject to $ax + by = c$, where $c > 0$ is some constant.

8. Let $f(x, y) = x^2(1 + y)^3 + 7y^2$ define a function on \mathbb{R}^2 . Find and classify its critical points. What is $\sup\{f(x, y) : (x, y) \in \mathbb{R}^2\}$? What is $\inf\{f(x, y) : (x, y) \in \mathbb{R}^2\}$?
9. Define $f(x) = (\log x)^{\log x}$, for $x > 1$. Using the chain rule, compute $f'(x)$.
10. Folland, §2.9, problem 16.
11. Suppose $F(x, y)$ is a C^2 function that satisfies the equations $F(x, y) = F(y, x)$, $F(x, x) = x$. Prove that the quadratic term in the Taylor polynomial of F based at the point (a, a) is $\frac{1}{2}F_{xx}(a, a)(x - y)^2$.
12. There may be homework problems or example problems from the text or lectures on the midterm.
13. The following topics have been covered since the first midterm:
 - (a) Chain rule.
 - (b) Mean value theorem.
 - (c) Higher order partials and equality of mixed partials.
 - (d) Taylor's theorem in one and several variables with Lagrange's form of the remainder.
 - (e) Behavior near critical points – second derivative test for extrema in the case of two variables.
 - (f) Max-min problems with constraints. The method of Lagrange multipliers.
 - (g) Implicit Function Theorem.
 - (h) Smooth curves and surfaces.
 - (i) Transformations.