

## Math 335 Sample Problems

One notebook-sized page of notes (both sides may be used) will be allowed on the final exam. No electronic devices allowed. The final will be comprehensive.

1. Prove that

$$\sum_{n=1}^{\infty} \frac{\cos(nx)}{n^{2/3}}$$

is not a Fourier series.

2. Suppose  $\int_a^{\infty} f(x)dx$  converges absolutely. Prove that

$$\lim_{s \rightarrow \pm\infty} \int_a^{\infty} f(x)e^{isx} dx = 0.$$

3. Fix  $r$  with  $0 \leq r < 1$  and consider the series

$$1 + 2re^{ix} + 3r^2e^{i2x} + 4r^3e^{i3x} + \dots$$

(a) Prove that this is a Fourier series.

(b) Prove that it converges uniformly on  $x \in \mathbb{R}$  and find the limit.

4. Suppose  $f$  is continuous and piecewise smooth. Prove that

$$\sum_{n \neq 0} |\hat{f}(n)| \leq \left( 2 \sum_1^{\infty} \frac{1}{n^2} \right)^{1/2} \frac{1}{\sqrt{2\pi}} \left( \int_{-\pi}^{\pi} |f'|^2 \right)^{1/2} = \sqrt{\frac{\pi}{6}} \left( \int_{-\pi}^{\pi} |f'|^2 \right)^{1/2}$$

5. Prove that

$$\int_0^1 (1-t^4)^{-1/2} dt = \frac{\Gamma(\frac{5}{4})\sqrt{\pi}}{\Gamma(\frac{3}{4})}.$$

6. Let  $f$  be a  $2\pi$ -periodic function and let  $a$  be a fixed real number and let a new function  $g$  be defined by  $g(x) = f(x-a)$ . What is the relation between the Fourier coefficients  $\hat{f}(n)$  and  $\hat{g}(n)$ ?
7. Find the Fourier series of the following function of  $x$ .

$$\frac{1-r^2}{1-2r\cos x+r^2}$$

where  $r$  is fixed with  $0 \leq r < 1$ . (You don't need to integrate.)

8. Let  $f$  be a  $2\pi$ -periodic, piecewise smooth function. Let  $\widehat{f}(n)$  be the complex Fourier coefficients of  $f$ . Show that there is a constant  $M$  (which will depend on  $f$ ) such that  $|\widehat{f}(n)| < M/|n|$  for all  $n \neq 0$ . Do **not** assume  $f$  is continuous.
9. Suppose  $f$  is Riemann integrable, and  $f_k$  is a sequence of Riemann integrable functions on  $[0, 2\pi]$  such that  $\lim_{k \rightarrow \infty} \int_0^{2\pi} |f_k - f| = 0$ . Prove that the Fourier coefficients satisfy  $\lim_{k \rightarrow \infty} \widehat{f}_k(n) = \widehat{f}(n)$  for each  $n$ .
10. Suppose  $f$  and  $g$  are  $2\pi$ -periodic and Riemann integrable on compact subsets of  $\mathbf{R}$ . Suppose also that  $f(x) = g(x)$  in a neighborhood of a point  $x_0$ . Suppose that the Fourier series for one of the functions converges at  $x_0$ . Prove that the other series converges at  $x_0$  and

$$\sum_{-\infty}^{\infty} \widehat{f}(n) e^{inx_0} = \sum_{-\infty}^{\infty} \widehat{g}(n) e^{inx_0}.$$

Hint: Look at  $f - g$ .

11. Prove that

$$\lim_{n \rightarrow \infty} \int_0^{\pi} \frac{\sin(nx)}{x} dx = \frac{\pi}{2}.$$

12. Define a function  $\log_p(x)$  inductively by the formulas  $\log_0(x) = x$ ,  $\log_{p+1}(x) = \log(\log_p(x))$ . Prove by induction that the series

$$\sum_{n=m}^{\infty} \frac{1}{\log_0(n) \log_1(n) \log_2(n) \dots \log_p(n)}$$

(where  $m$  is large enough for the denominators to be defined as real numbers) diverges for every  $p$ .

13. Suppose that  $a_n > 0$ , that  $a_n$  is decreasing, and that  $\sum_1^{\infty} a_n$  converges. Is it true that  $\lim_{n \rightarrow \infty} na_n = 0$ ? If true prove it, if false give a counterexample.

14. Suppose that  $f$  is  $2\pi$ -periodic, continuous, and piecewise linear (that means that there is a finite set (in  $[-\pi, \pi]$ ) of intervals in each of which  $f$  is defined by a linear function). Prove that

$$|\widehat{f}(n)| \leq \frac{c}{n^2},$$

for some constant  $c$ .

15. Show that the series  $\sum_1^{\infty} \frac{\sin nx}{\sqrt{n}}$  converges for all  $x$  and uniformly on any interval of the form  $[\delta, 2\pi - \delta]$ , where  $\delta > 0$  is small. Show that the series is not the Fourier series of a Riemann integrable function.
16. Find the solution of  $u_t = 3u_{xx}$ ,  $u(0, t) = u(\pi, t) = 0$ ,  $u(x, 0) = \cos x \sin 5x$ . (This is easier than it looks.)

17. (a) Let  $\sum_{n=0}^{\infty} a_n x^n$  be a series with radius of convergence  $R$ . Substitute  $re^{i\theta}$  for  $x$  and get a new series involving  $e^{in\theta}$ . If  $0 < r < R$  prove that this is a Fourier series (the variable is  $\theta$ ).
- (b) Prove that  $\sum_{n=0}^{\infty} r^{2n} |a_n|^2$  converges for  $0 \leq r < R$ .

18. Compute

$$\lim_{n \rightarrow \infty} \int_a^b \sin^2(nx) dx.$$

19. Folland, §8.6: problem 10.

20. Let  $f$  and  $g$  be continuous  $2\pi$ -periodic functions. Define the *convolution* of  $f$  and  $g$  to be the function.  
 $f * g(x) = \frac{1}{2\pi} \int_0^{2\pi} f(x-t)g(t)dt.$

- (a) Prove that  $f * g$  is  $2\pi$ -periodic.
- (b) Prove that  $\widehat{f * g}(n) = \widehat{f}(n)\widehat{g}(n)$ , so the Fourier series of  $f * g$  is  $\sum_{-\infty}^{\infty} c_n d_n e^{inx}$ , where  $c_n = \widehat{f}(n)$ ,  $d_n = \widehat{g}(n)$ .

21. (a) Find the cosine series of  $f$  where  
 $f(x) = 0, 0 < x < \pi/2; f(x) = 1, \pi/2 < x < \pi.$
- (b) Prove that the series converges for all  $x$ .
- (c) For which  $x$  does the series converge absolutely?

22. Suppose  $a_n > 0$  and  $\sum_{n=1}^{\infty} a_n$  converges. Let  $t_n = \sum_{k \geq n} a_k$ .

- (a) Prove that  $\sum \frac{a_n}{t_n}$  diverges.
- (b) Prove that  $\sum \frac{a_n}{\sqrt{t_n}}$  converges.

23. Suppose  $a_n > 0, b_n > 0$  suppose  $f(x) = \sum a_n x^n$  and  $g(x) = \sum b_n x^n$  converge for all  $x$ . Suppose also that  $\lim_{n \rightarrow \infty} \frac{a_n}{b_n} = c$ . Prove that

$$\lim_{x \rightarrow +\infty} \frac{f(x)}{g(x)} = c.$$

24. (a) Let  $r = \sqrt{x^2 + y^2}$ . Prove that  $\frac{y}{r^2}$  is harmonic when  $y > 0$ .
- (b) Suppose  $\phi(t)$  is continuous on  $[a, b]$ . Let

$$u(x, y) = \int_a^b \frac{\phi(t)y dt}{(x-t)^2 + y^2}.$$

Prove that  $u$  is harmonic when  $y > 0$ .

25. Suppose  $f(x)$  is  $2\pi$  periodic and satisfies  $|f(x) - f(y)| \leq M|x - y|$  for all  $x, y$ . Let

$$u(r, \theta) = \int_0^{2\pi} f(\theta + \phi) P_r(\phi) d\phi,$$

for  $0 < r < 1$ , where  $P_r(\phi)$  is the Poisson kernel. Prove that  $\frac{\partial u}{\partial \theta}$  exists and  $|\frac{\partial u}{\partial \theta}| \leq M$ .

26. let  $f$  be  $2\pi$ -periodic, continuous, and piecewise smooth. Let  $m$  be any positive integer and define the function  $f_m$  by the formula  $f_m(x) = f(mx)$ . Prove that  $\widehat{f_m}(n) = \widehat{f}\left(\frac{n}{m}\right)$  if  $m$  divides  $n$  and is 0 otherwise.

27. Determine  $a, b, c$  so that  $f_0(x) = 1, f_1(x) = x + a, f_2(x) = x^2 + bx + c$  is an orthogonal set using the inner product  $\langle f, g \rangle = \int_{-1}^1 fg$  on  $[-1, 1]$ .

28. (Extra, extra credit) Let  $(x)$  be the function with period 1 that equals  $x$  on  $(-1/2, 1/2)$  and equals 0 at  $\pm 1/2$ . Define a function  $f$  as follows

$$f(x) = \sum_1^{\infty} \frac{(nx)}{n^2}.$$

This is an example of Riemann.

- (a) Prove that the series defining (1) converges uniformly on  $\mathbb{R}$ .

- (b) Prove that  $f$  is continuous at points not of the form  $\frac{2s+1}{2n}$ .

- (c) If  $2s+1$  and  $n$  are relatively prime let  $a = \frac{2s+1}{2n}$ . Prove that the jump discontinuity at  $a$  is

$$\lim_{x \rightarrow a^+} f(x) - \lim_{x \rightarrow a^-} f(x) = -\frac{\pi^2}{8n^2} \text{ at } \frac{2s+1}{2n}.$$

- (d) Prove that  $f$  is Riemann integrable on each compact subinterval of  $\mathbb{R}$ .

29. There may be problems from the text, statements of theorems from the text, problems from previous review sets, or examples from class on the exam.