THE CUTSET-RANK COROLLARY

JOEL NISHIMURA, PETER MANNISTO

ABSTRACT. This paper contains a generalization of the relationship outlined in [1] between the connections of an electrical network Γ and the determinants of $\Lambda(P;Q)$. Specifically, it will be shown that rank $\Lambda(P;Q) \leq |M|$ where M is the minimal cutset of the graph of Γ . In addition, when the graph is circular planar, rank $\Lambda(P;Q) = |M|$.

Let $\Gamma=(G,\gamma)$ be an electrical network, and let the vertices V of G be partitioned into $V=\partial V\cup$ int V, where $\partial V\neq\emptyset$. The response matrix Λ of Γ is defined as the matrix for which $\Lambda\phi=\psi$, where ϕ is the column vector of voltages defined on ∂V , and ψ is the vector of currents on ∂V . Let P and Q be two disjoint subsets of ∂V , where |P|=|Q|. Then $\Lambda(P;Q)$ is the submatrix of Λ with row indices P and column indices Q. In [1] it is proved that if det $\Lambda(P;Q)\neq 0$ then there exists a k-connection from P to Q, where k=|P|=|Q|. Moreover, in the circular planar case the existence of a k-connection implies that det $\Lambda(P;Q)\neq 0$. Our generalization of these two statements will require the following definitions:

Definition 1. (Cutset, Minimal Cutset): Given a graph $G = \{V, E\}$ (where V is the set of vertices and E is the set of edges) and two disjoint sets of vertices $P, Q \subset V$ a cutset of the pair P, Q is a set $R \subset V$ such that all paths from a vertex $P \in P$ to a vertex $P \in Q$ intersect P. A minimal cutset of the pair P, P is a cutset of P, P of minimal cardinality among all such cutsets.

We can now state Menger's Theorem:

Theorem 1. (Menger's Theorem [2]): Let $G = \{V, E\}$ be a graph and $A, B \subseteq V$. Then the cardinality of the minimum cutset of the pair A, B in G is equal to the maximum number of disjoint paths from A to B in G.

Definition 2. (Degree of Connectivity) Given two disjoint subsets of ∂V P and Q, the degree of connectivity of (P,Q) is defined as the largest integer k such that there exists a k-connection between P' and Q', where $P' \subseteq P$ and $Q' \subseteq Q$.

Our proof of the cutset-rank corrolary requires the following lemma.

Lemma 1. Given a graph G, define a graph G' where $V' = V - \partial V + P + Q$ and $E' = E - E(\partial V)$ where $E(\partial V)$ are all edges incident on a deleted vertex. On this graph G', the degree of connectivity is equal to the cardinality of the minimum cutset.

This follows directly from Menger's Theorem as the degree of connectivity is precisely the maximum number of disjoint paths from P to Q.

We now head on to the desired result.

Date: July 26, 2005.

Corollary 1. (Cutset-Rank Corollary) Given $\Lambda(P;Q)$ defined on $\Gamma=(G,\gamma)$. Let the minimal cutset of (P,Q) be denoted by M. In general,

(1)
$$\operatorname{rank} \Lambda(P; Q) \le |M|$$

Moreover, if G is circular planar,

(2)
$$\operatorname{rank} \Lambda(P; Q) = |M|$$

Proof. Let $\Lambda(P;Q)$ have rank m. This implies that the largest non-zero sub-determinant of $\Lambda(P;Q)$ is of size $m \times m$. Therefore there exists an m-connection between $P' \subseteq P$ and $Q' \subseteq Q$ where |Q'| = |P'|. Thus the degree of connectivity is at least m, the rank of $\Lambda(P;Q)$.

Moreover, if G is circular planar, since every subdeterminant larger than $m \times m$ of $\Lambda(P;Q)$ is zero then there does not exist any k-connection for k > m. This implies then that for the circular planar case the degree of connectivity is equal to m, the rank of $\Lambda(P;Q)$.

By Lemma 1 both of these statements can translated into the desired results. Q.E.D.

References

- [1] Curtis, Edward B. & Morrow, James A., *Inverse Problems for Electrical Networks*, Series on Applied Mathematics Vol. 13, World Scientifc, New Jersey, 2000.
 - [2] Diestel, Richard, Graph Theory, Springer-Verlag, New York, 1997.