

MATH 112 - Winter 2005
Exam 2, Version 2 - Hints and Answers

1. (4 points each)

(a) $f'(t) = (t^4 - 6t^3 + 4t)(5t^4 + 28t^3 + 4t) + (t^5 + 7t^4 + 2t^2)(4t^3 - 18t^2 + 4)$

(b) $R'(q) = \frac{(1 + 3e^q) \cdot \frac{1}{q^2 - 5q} \cdot (2q - 5) - \ln(q^2 - 5q) \cdot (3e^q)}{(1 + 3e^q)^2}$

(c) $f_x(x, y) = 2xe^y + (e^{x^2})(2x)y + e^{x^2y}(2xy)$

(d) $f_y(x, y) = x^2e^y + e^{x^2} + e^{x^2y}(x^2)$

2. (10 points) HINT: The five vertices are (0, 2), (0, 10), (5, 9), (7, 0) and (3, 0).

ANSWER: maximum=87; minimum=9

3. (4 points each)

(a) HINT: $MC(q) = 0.03q^2 - 0.6q + 3$. The question asks where this function has a horizontal tangent line. So, you must compute the q at which $MC'(q) = 0$. $MC'(q) = 0.06q - 0.6$. Set this equal to 0 and solve for t .

ANSWER: $q = 10$

(b) i. ANSWER: global maximum= $MC(14) = 0.48$; global minimum= $MC(11) = 0.03$

ii. ANSWER: global maximum= $MC(6) = 0.48$; global minimum= $MC(10) = 0$

(c) ANSWER: $P(q) = -0.01q^3 + 0.22q^2 - 0.65q - 4$

$P'(q) = -0.03q^2 + 0.44q - 0.65$. We need to know if P has a horizontal tangent line at $q = 13$. Plugging 13 into P' gives $P'(13) = 0$. So, P has a horizontal tangent line at $q = 13$.

$P''(q) = -0.06q + 0.44$ Plugging 13 into P'' gives $P''(13) = -0.34$, which is negative. So, P is concave down at 13.

Since P has a horizontal tangent and is concave down at $q = 13$, $q = 13$ gives a local maximum of profit.

4. (4 points each)

(a) HINT: $f_x(x, y) = x - 5 + 2y$ and $f_y(x, y) = -y - 4 + 2x$. Set these partial derivatives equal to 0 and solve the resulting system of equations.

ANSWER: $(x, y) = (2.6, 1.2)$

(b) HINT: $A \approx f_y(4, 4) = 0$; $B \approx f_x(4, 4) = 7$

ANSWER: B is bigger

(c) HINT: $h'(1) = f_y(6, 1) = 7$, $k'(5) = f_x(5, 2) = 4$

ANSWER: i is steeper