


# Jordan Content

Note Title

11/21/2009

Theorem: A bounded set  $S$  is Jordan measurable if and only if the outer area of  $\partial S$  is 0.

Proof:  $S$  is measurable exactly when  $\chi_S$  is Riemann integrable. I will omit the proof that integrability is independent of the rectangle containing  $S$  (this does require proof). So assume



$\overline{S}$  is in the interior of the containing rectangle,

Suppose  $\chi_S$  is integrable. Then there is a partition

so that  $S_p(\chi_S) - s_p(\chi_S) < \epsilon$ . Let



$T = \{R_{ij} : S \cap R_{ij} \neq \emptyset, \text{ and } S^c \cap R_{ij} \neq \emptyset\}$ . Then

$$S_p(\chi_S) - s_p(\chi_S) = \sum_{R_{ij} \in T} |R_{ij}| < \epsilon.$$

Let  $D = \bigcup_{R_{ij} \in T} R_{ij}$ . I claim that  $\partial S \subset D$ .

Let  $p \in \partial S$ . Then if  $p \in \text{int}(R_{ij})$ ,  $S \cap R_{ij} \neq \emptyset$  and  $S^c \cap R_{ij} \neq \emptyset$ . So  $R_{ij} \in T$ . If  $p \in \partial R_{ij}$ , then

$p$  is a corner or edge of  $R_{ij}$  and we have one of the following figures




Then for one of the adjacent rectangles  $R_j$  it must be true that there is a point of  $S$  and a point of  $S^c$ . Since  $p$  is in all of these rectangles,  $p \in R_{ij}$  for some  $R_{ij} \in T$ . This proves  $\partial S \subset D$ .

Since  $\sum |R_{ij}| < \epsilon$ ,  $\partial S$  has content. may contribute

Next suppose  $\bar{A}(\partial S) = 0$ .

We have a finite union of rectangles  $R_\alpha$  with sides



parallel to the axes and  $\sum |R_\alpha| < \epsilon$ . We can create a partition so that these rectangles are unions of

rectangles in the partition (see dotted lines for examples). Unfortunately it may not happen that


$$S_p(x_S) - A_p(x_S) = \sum |R_\alpha|. \quad (\text{The brown rectangle})$$

may have points of  $S$  and  $S^c$ .)

However by adding additional lines we can create a refinement of the partition and with this partition

$$S_p(\chi_S) - L_p(\chi_S) < 2\epsilon, \text{ say. So } \chi_S \text{ is integrable.}$$

Or we could replace the original rectangles  $R_\alpha$  with slightly larger rectangles  $R'_\alpha$ , whose interiors cover  $\partial S$  ( $\partial S$  is compact) and then the other rectangles have no points of  $\partial S$ . Since rectangles are convex, these other rectangles are entirely contained in  $S$  or  $S^c$ .



Why doesn't this proof work to show that an integrable function is continuous except on a set of Jordan content 0 (which is not true)?