

Math 334 Sample Problems

One side of one notebook sized page of notes will be allowed on the test. You may work together on the sample problems – I encourage you to do that. The test will cover up to §4.3.

1. Expand $(1 - x + 2y)^3$ in powers of $x - 1$ and $y - 2$ in two different ways. The first way is by using algebra and the second way is by computing the Taylor series.
2. Let f be defined and bounded on $[a, b]$. Define a function g on $[a, b]$ by the formula $g(x) = \bar{I}(\chi_{[a,x]}f)$. In other words $g(x)$ is the upper integral of f on the interval $[a, x]$. Prove that g is continuous on $[a, b]$. Suppose f is continuous at x_0 . Prove that $g'(x_0) = f(x_0)$. The same is true for lower integrals.
3. Folland, §4.1, # 9.
4. Suppose $a < b < c < d$. Let $I = [a, b]$, $J = [c, d]$, $R = I \times J$ and let $f(x, y) = |x - y|$, if $x \in I$, $y \in J$. Compute $\int_R f$.
5. Using the method of Lagrange multipliers, find the highest and lowest points of the circle
$$x^2 + y^2 + z^2 = 16, (x + 1)^2 + (y + 1)^2 + (z + 1)^2 = 27$$
6. Show that the surface $z = 3x^2 - 2xy + 2y^2$ lies entirely above every one of its tangent planes. Hint: Look at the Taylor expansion at every point.

7. Find the shortest distance from the point $(1, -1, 1)$ to the surface $z = xy$.
8. Let f be continuously differentiable on $[a, b]$ and assume $f'(x) > 0$ on $[a, b]$.
- (a) Prove that f has a continuously differentiable inverse g and that $g'(x) > 0$.
- (b) Prove that

$$\int_a^b f + \int_{f(a)}^{f(b)} g = bf(b) - af(a).$$

Can you give a geometric interpretation of the result?

9. Let $f(x, y) = \sec(x + y^2)$. Find the first two non-zero terms in the Taylor series of $\cos x$, centered at 0. Use it to find the first two non-zero terms of the Taylor series of $\sec x$ centered at 0. Then use that series to find the first two non-zero terms of f at $(0, 0)$.
10. Let g be a polynomial of degree three. Prove that

$$\int_{-1}^1 g = \frac{g(-1) + 4g(0) + g(1)}{3}.$$

11. Consider the following function

$$F(x, y) = \left(\frac{x}{1 + x + y}, \frac{y}{1 + x + y} \right),$$

which has the set $\{(x, y) : 1 + x + y \neq 0\}$ as its domain. Compute $\frac{\partial(f, g)}{\partial(x, y)}$. Where is it different from 0? Show that F is 1 - 1 and find an explicit formula for its inverse. Use these results to describe the exact image of F

12. Folland, §2.9, problem 16.

13. Let f be a positive continuous function on $I = [a, b]$. Let $M = \max\{f(x) : x \in I\}$. Prove that

$$\lim_{n \rightarrow \infty} \left(\int_I f^n \right)^{1/n} = M.$$

14. Suppose $F(x, y)$ is a C^2 function that satisfies the equations $F(x, y) = F(y, x)$, $F(x, x) = x$. Prove that the quadratic term in the Taylor polynomial of F based at the point (a, a) is $\frac{1}{2}F_{xx}(a, a)(x - y)^2$.
15. There may be homework problems or example problems from the text or lectures on the midterm.